

PLC η 1 siRNA (h): sc-77989

BACKGROUND

Phosphoinositide-specific phospholipase C (PLC) plays a crucial role in the initiation of receptor mediated signal transduction through the generation of the two second messengers, inositol 1,4,5-triphosphate and diacylglycerol from phosphatidylinositol 4,5-bisphosphate. PLC isozymes are divided into subclasses based on structure and activation mechanisms. PLC η 1 (1-phosphatidylinositol-4,5-bisphosphate phosphodiesterase eta-1), also known as Phospholipase C-like protein 3 (PLC-L3) is a 1693 amino acid protein that is especially found in areas of the cerebellum, spinal cord and cerebrum. PLC η 1 contains a C2 domain, a PH domain, a PI-PLC X-box domain, a PI-PLC Y-box domain and two EF-hand domains through which it binds calcium to serve as a cofactor. There are four isoforms of PLC η 1 that are produced as a result of alternative splicing events.

REFERENCES

1. Rhee, S.G. 2001. Regulation of phosphoinositide-specific phospholipase C. *Annu. Rev. Biochem.* 70: 281-312.
2. Hwang, J.I., et al. 2005. Molecular cloning and characterization of a novel phospholipase C, PLC- η . *Biochem. J.* 389: 181-186.
3. Katan, M. 2005. New insights into the families of PLC enzymes: looking back and going forward. *Biochem. J.* 391: e7-e9.
4. Stewart, A.J., et al. 2005. Identification of a novel class of mammalian phosphoinositide-specific phospholipase C enzymes. *Int. J. Mol. Med.* 15: 117-121.
5. Nakahara, M., et al. 2005. A novel phospholipase C, PLC η 2, is a neuron-specific isozyme. *J. Biol. Chem.* 280: 29128-29134.
6. Stewart, A.J., et al. 2007. Phospholipase C- η enzymes as putative protein kinase C and Ca^{2+} signalling components in neuronal and neuroendocrine tissues. *Neuroendocrinology* 86: 243-248.
7. Imai, T., et al. 2007. Expression and characterization of a pleckstrin homology domain in phospholipase C, PLC- η 1. *Protein Expr. Purif.* 56: 247-252.
8. Suh, P.G., et al. 2008. Multiple roles of phosphoinositide-specific phospholipase C isozymes. *BMB Rep.* 41: 415-434.

CHROMOSOMAL LOCATION

Genetic locus: PLCH1 (human) mapping to 3q25.31.

PRODUCT

PLC η 1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PLC η 1 shRNA Plasmid (h): sc-77989-SH and PLC η 1 shRNA (h) Lentiviral Particles: sc-77989-V as alternate gene silencing products.

For independent verification of PLC η 1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-77989A, sc-77989B and sc-77989C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PLC η 1 siRNA (h) is recommended for the inhibition of PLC η 1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PLC η 1 gene expression knockdown using RT-PCR Primer: PLC η 1 (h)-PR: sc-77989-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.