

# SPATA6 siRNA (h): sc-78827

## BACKGROUND

SPATA6 (spermatogenesis associated 6), also known as SRF1, is a 488 amino acid secreted protein that may be involved in spermatid maturation or sperm function. SPATA6 is expressed during embryonic development and is localized in neural tube, somites and limb buds of mouse embryo. Existing as two isoforms produced by alternative splicing events, the gene encoding SPATA6 maps to mouse chromosome 4 D1 and human chromosome 1p33. Human chromosome 1 spans 260 million base pairs, contains over 3,000 genes and comprises nearly 8% of the human genome. Chromosome 1 houses a large number of disease-associated genes, including those that are involved in familial adenomatous polyposis, Stickler syndrome, Parkinson's disease, Gaucher disease, schizophrenia and Usher syndrome. Aberrations in chromosome 1 are found in a variety of cancers, including head and neck cancer, malignant melanoma and multiple myeloma.

## REFERENCES

1. Watson, M.L., et al. 1990. Genomic organization of the selectin family of leukocyte adhesion molecules on human and mouse chromosome 1. *J. Exp. Med.* 172: 263-272.
2. Onisto, M., et al. 2001. Evidence for FSH-dependent upregulation of SPATA2 (spermatogenesis-associated protein 2). *Biochem. Biophys. Res. Commun.* 283: 86-92.
3. Blackwood, D.H., et al. 2001. Schizophrenia and affective disorders— cosegregation with a translocation at chromosome 1q42 that directly disrupts brain-expressed genes: clinical and P300 findings in a family. *Am. J. Hum. Genet.* 69: 428-433.
4. Liliana Slongo, M., et al. 2003. Cloning and characterization of the promoter region of human SPATA2 (spermatogenesis-associated protein 2) gene. *Biochim. Biophys. Acta* 1625: 192-196.
5. Oh, C., et al. 2003. Characterization, expression pattern and chromosomal localization of the spermatogenesis associated 6 gene (Spata6). *Mol. Hum. Reprod.* 9: 321-330.
6. Weise, A., et al. 2005. New insights into the evolution of chromosome 1. *Cytogenet. Genome Res.* 108: 217-222.
7. Deng, Y., et al. 2006. Expression and identification of a novel apoptosis gene Spata17 (MSRG-11) in mouse spermatogenic cells. *Acta Biochim. Biophys. Sin.* 38: 37-45.
8. Gregory, S.G., et al. 2006. The DNA sequence and biological annotation of human chromosome 1. *Nature* 441: 315-321.
9. Moro, E., et al. 2007. Zebrafish SPATA2 is expressed at early developmental stages. *Int. J. Dev. Biol.* 51: 241-246.

## CHROMOSOMAL LOCATION

Genetic locus: SPATA6 (human) mapping to 1p33.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## PRODUCT

SPATA6 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SPATA6 shRNA Plasmid (h): sc-78827-SH and SPATA6 shRNA (h) Lentiviral Particles: sc-78827-V as alternate gene silencing products.

For independent verification of SPATA6 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-78827A, sc-78827B and sc-78827C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

SPATA6 siRNA (h) is recommended for the inhibition of SPATA6 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SPATA6 gene expression knockdown using RT-PCR Primer: SPATA6 (h)-PR: sc-78827-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.