

# TLR1 (GD2.F4): sc-80535

## BACKGROUND

Six human homologs of the *Drosophila* Toll receptor were initially identified based on their sequence similarities and designated Toll-like receptors (TLR). Toll receptors are involved in mediating dorsoventral polarization in the developing *Drosophila* embryo and also participate in the host immunity. The TLR family of proteins are characterized by a highly conserved Toll homology (TH) domain, which is essential for Toll-induced signal transduction. TLR1, as well as the other TLR family members, is a type I transmembrane receptor that characteristically contains an extracellular domain consisting of several leucine-rich regions along with a single cytoplasmic Toll/IL-1R-like domain. TLR2 and TLR4 are activated in response to lipopolysaccharide (LPS) stimulation, which results in the activation and translocation of NF $\kappa$ B and suggests that these receptors are involved in mediating inflammatory responses. Expression of TLR receptors is highest in peripheral blood leukocytes, macrophages and monocytes. TLR6 is highly homologous to TLR1, sharing greater than 65% sequence identity and, like other members of the TLR family, it induces NF $\kappa$ B signaling upon activation.

## REFERENCES

1. Gay, N.J. and Keith, F.J. 1991. *Drosophila* Toll and IL-1 receptor. *Nature* 351: 355-356.
2. Medzhitov, R., Preston-Hurlburt, P. and Janeway, C.A., Jr. 1997. A human homologue of the *Drosophila* Toll protein signals activation of adaptive immunity. *Nature* 388: 394-397.
3. Rock, F.L., Hardiman, G., Timans, J.C., Kastelein, R.A. and Bazan, J.F. 1998. A family of human receptors structurally related to *Drosophila* Toll. *Proc. Natl. Acad. Sci. USA* 95: 588-593.
4. Yang, R.B., Mark, M.R., Gray, A., Huang, A., Xie, M.H., Zhang, M., Goddard, A., Wood, W.I., Gurney, A.L. and Godowski, P.J. 1998. Toll-like receptor-2 mediates lipopolysaccharide-induced cellular signalling. *Nature* 395: 284-288.
5. Brightbill, H.D., Libraty, D.H., Krutzik, S.R., Yang, R.B., Belisle, J.T., Bleharski, J.R., Maitland, M., Norgard, M.V., Plevy, S.E., Smale, S.T., Brennan, P.J., Bloom, B.R., Godowski, P.J. and Modlin, R.L. 1999. Host defense mechanisms triggered by microbial lipoproteins through Toll-like receptors. *Science* 285: 732-736.
6. Chow, J.C., Young, D.W., Golenbock, D.T., Christ, W.J. and Gusovsky, F. 1999. Toll-like receptor-4 mediates lipopolysaccharide-induced signal transduction. *J. Biol. Chem.* 274: 10689-10692.
7. Schwandner, R., Dziarski, R., Wesche, H., Rothe, M. and Kirschning, C.J. 1999. Peptidoglycan- and lipoteichoic acid-induced cell activation is mediated by Toll-like receptor 2. *J. Biol. Chem.* 274: 17406-17409.
8. Takeuchi, O., Kawai, T., Sanjo, H., Copeland, N.G., Gilbert, D.J., Jenkins, N.A., Takeda, K. and Akira, S. 1999. TLR6: A novel member of an expanding Toll-like receptor family. *Gene* 231: 59-65.

## CHROMOSOMAL LOCATION

Genetic locus: TLR1 (human) mapping to 4p14.

## SOURCE

TLR1 (GD2.F4) is a mouse monoclonal antibody raised against human recombinant soluble TLR1 extracellular domain (TLR1-Fc).

## PRODUCT

Each vial contains 200  $\mu$ g IgG<sub>1</sub> kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

TLR1 (GD2.F4) is available conjugated to either phycoerythrin (sc-80535 PE) or fluorescein (sc-80535 FITC), 200  $\mu$ g/ml, for IF, IHC(P) and FCM.

## APPLICATIONS

TLR1 (GD2.F4) is recommended for detection of TLR1 of human origin by immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500) and flow cytometry (1  $\mu$ g per  $1 \times 10^6$  cells).

Suitable for use as control antibody for TLR1 siRNA (h): sc-40254, TLR1 shRNA Plasmid (h): sc-40254-SH and TLR1 shRNA (h) Lentiviral Particles: sc-40254-V.

Molecular Weight of TLR1: 90 kDa.

## RECOMMENDED SUPPORT REAGENTS

To ensure optimal results, the following support reagents are recommended:  
 1) Immunofluorescence: use m-IgG $\kappa$  BP-FITC: sc-516140 or m-IgG $\kappa$  BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz<sup>®</sup> Mounting Medium: sc-24941 or UltraCruz<sup>®</sup> Hard-set Mounting Medium: sc-359850.

## STORAGE

Store at 4<sup>°</sup> C, \*\*DO NOT FREEZE\*\*. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.