

p-MEK-1/2 (7E10): sc-81503

BACKGROUND

A family of protein kinases located upstream of the MAP kinases and responsible for their activation has been identified. The prototype member of this family, designated MAP kinase kinase, or MEK-1, specifically phosphorylates the MAP kinase regulatory threonine and tyrosine residues present in the Thr-Glu-Tyr motif of ERK. A second MEK family member, MEK-2, resembles MEK-1 in its substrate specificity. MEK-3 (or MKK-3) functions to activate p38 MAP kinase, and MEK-4 (also called SEK1 or MKK-4) activates both p38 and JNK MAP kinases. MEK-5 appears to specifically phosphorylate ERK5, whereas MEK-6 phosphorylates p38 and p38 β . MEK-7 (or MKK-7) phosphorylates and activates the JNK signal transduction pathway. Phosphorylation on Ser/Thr by MAP kinase kinase kinases (RAFor MEKK1) positively regulates the kinase activity.

REFERENCES

1. Crews, C.M., et al. 1992. The primary structure of MEK, a protein kinase that phosphorylates the ERK gene product. *Science* 258: 478-480.
2. Wu, J., et al. 1993. Identification and characterization of a new mammalian mitogen-activated protein kinase kinase, MKK-2. *Mol. Cell. Biol.* 13: 4539-4548.
3. Dérijard, B., et al. 1995. Independent human MAP-kinase signal transduction pathways defined by MEK and MKK isoforms. *Science* 267: 682-685.
4. Zhou, G., et al. 1995. Components of a new human protein kinase signal transduction pathway. *J. Biol. Chem.* 270: 12665-12669.

CHROMOSOMAL LOCATION

Genetic locus: MAP2K1 (human) mapping to 15q22.31, MAP2K2 (human) mapping to 19p13.3; Map2k1 (mouse) mapping to 9 C, Map2k2 (mouse) mapping to 10 C1.

SOURCE

p-MEK-1/2 (7E10) is a mouse monoclonal antibody raised against phosphopeptide corresponding to amino acid residues surrounding the T-E-Y motif of MEK-1/2 of human origin.

PRODUCT

Each vial contains 50 μ g IgG₁ in 0.5 ml of PBS with < 0.1% sodium azide, 0.1% gelatin, PEG and sucrose.

APPLICATIONS

p-MEK-1/2 (7E10) is recommended for detection of Ser 218 and 222 phosphorylated MEK-1 and Ser 222 and 226 phosphorylated MEK-2 of mouse, rat, human and canine origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) and immunoprecipitation [1-2 μ g per 100-500 μ g of total protein (1 ml of cell lysate)].

Molecular Weight of MEK-1: 45 kDa.


Molecular Weight of MEK-2: 47 kDa.

Positive Controls: A549 cell lysate: sc-2413, HeLa-PMA cell lysate: sc-2258 or NIH/3T3 whole cell lysate: sc-2210.


STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



Western blot analysis of MEK-1/2 phosphorylation in untreated (A, D), PDGF treated (B, E) and PDGF and lambda protein phosphatase (sc-200312A) treated (C, F) NIH/3T3 whole cell lysates. Antibodies tested include p-MEK-1/2 (7E10): sc-81503 (A, B, C) and MEK-1 (H-8): sc-6250 (D, E, F).



Western blot analysis of MEK-1/2 phosphorylation in untreated (A, D), serum-starved, PMA-treated (B, E), serum-starved, PMA and lambda protein phosphatase (sc-200312A) treated (C, F) HeLa whole cell lysates. Antibodies tested include p-MEK-1/2 (7E10): sc-81503 (A, B, C) and MEK-1 (H-8): sc-6250 (D, E, F).

SELECT PRODUCT CITATIONS

1. Kumarasamy, V.M., et al. 2015. Selective repression of RET proto-oncogene in medullary thyroid carcinoma by a natural alkaloid berberine. *BMC Cancer* 15: 599.
2. Zhang, D., et al. 2017. Downregulation of ATP1A1 promotes cancer development in renal cell carcinoma. *Clin. Proteomics* 14: 15.
3. Kim, I.H. and Nam, T.J. 2018. Fucoidan downregulates Insulin-like growth factor-I receptor levels in HT-29 human colon cancer cells. *Oncol. Rep.* 39: 1516-1522.
4. Kang, Y.M., et al. 2018. Inhibitory effects of bee venom on mast cell-mediated allergic inflammatory responses. *Int. J. Mol. Med.* 41: 3717-3726.
5. Liao, G., et al. 2018. MicroRNA miR-509 regulates ERK-1/2, the Vimentin network, and focal adhesions by targeting Plk1. *Sci. Rep.* 8: 12635.
6. Liu, P., et al. 2019. Crude protein from spirulina increases the viability of CCD-986sk cells via the EGFR/MAPK signaling pathway. *Int. J. Mol. Med.* 43: 771-778.
7. Kano, Y., et al. 2019. Tyrosyl phosphorylation of KRAS stalls GTPase cycle via alteration of switch I and II conformation. *Nat. Commun.* 10: 224.
8. Xing, J., et al. 2019. TRPM7 channel inhibition exacerbates pulmonary arterial hypertension through MEK/ERK pathway. *Aging* 11: 4050-4065.
9. Wang, J., et al. 2019. FDA-approved drug screen identifies proteasome as a synthetic lethal target in MYC-driven neuroblastoma. *Oncogene* 38: 6737-6751.
10. Chen, H., et al. 2020. TGF- β 1/IL-11/MEK/ERK signaling mediates senescence-associated pulmonary fibrosis in a stress-induced premature senescence model of Bmi-1 deficiency. *Exp. Mol. Med.* 52: 130-151.

RESEARCH USE

For research use only, not for use in diagnostic procedures.