

p-MEK-3/6 (B-9): sc-8407



The Power to Question

BACKGROUND

A family of protein kinases located upstream of the MAP kinases and responsible for their activation has been identified. The prototype member of this family, designated MAP kinase kinase, or MEK-1, specifically phosphorylates the MAP kinase regulatory threonine and tyrosine residues present in the Thr-Glu-Tyr motif of ERK. A second MEK family member, MEK-2, resembles MEK-1 in its substrate specificity. MEK-3 (or MKK-3) functions to activate p38 MAP kinase, and MEK-4 (also called SEK1 or MKK-4) activates both p38 and JNK MAP kinases. MEK-5 appears to specifically phosphorylate ERK5, whereas MEK-6 phosphorylates p38 and p38b. MEK-7 (or MKK-7) phosphorylates and activates the JNK signal transduction pathway.

CHROMOSOMAL LOCATION

Genetic locus: MAP2K3 (human) mapping to 17q11.2, MAP2K6 (human) mapping to 17q24.3; Map2k3 (mouse) mapping to 11 B2, Map2k6 (mouse) mapping to 11 E2.

SOURCE

p-MEK-3/6 (B-9) is a mouse monoclonal antibody raised against a sequence containing Ser 189 and Ser 207 phosphorylated MEK-3/6 of human origin.

PRODUCT

Each vial contains 200 µg IgG₁ kappa light chain in 1.0 ml of PBS with < 0.1% sodium azide and 0.1% gelatin.

p-MEK-3/6 (B-9) is available conjugated to agarose (sc-8407 AC), 500 µg/0.25 ml agarose in 1 ml, for IP; to HRP (sc-8407 HRP), 200 µg/ml, for WB, IHC(P) and ELISA; to either phycoerythrin (sc-8407 PE), fluorescein (sc-8407 FITC), Alexa Fluor® 488 (sc-8407 AF488), Alexa Fluor® 546 (sc-8407 AF546), Alexa Fluor® 594 (sc-8407 AF594) or Alexa Fluor® 647 (sc-8407 AF647), 200 µg/ml, for WB (RGB), IF, IHC(P) and FCM; and to either Alexa Fluor® 680 (sc-8407 AF680) or Alexa Fluor® 790 (sc-8407 AF790), 200 µg/ml, for Near-Infrared (NIR) WB, IF and FCM.

Blocking peptide available for competition studies, sc-8407 P, (100 µg peptide in 0.5 ml PBS containing < 0.1% sodium azide and 0.2% stabilizer protein).

Alexa Fluor® is a trademark of Molecular Probes, Inc., Oregon, USA

APPLICATIONS

p-MEK-3/6 (B-9) is recommended for detection of MEK-3 and MEK-6 phosphorylated at Ser 189 and Ser 207 of mouse, rat and human origin by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000), immunoprecipitation [1-2 µg per 100-500 µg of total protein (1 ml of cell lysate)], immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500), immunohistochemistry (including paraffin-embedded sections) (starting dilution 1:50, dilution range 1:50-1:500) and solid phase ELISA (starting dilution 1:30, dilution range 1:30-1:3000).


Suitable for use as control antibody for MEK-3/6 siRNA (h): sc-43924, MEK-3/6 shRNA Plasmid (h): sc-43924-SH and MEK-3/6 shRNA (h) Lentiviral Particles: sc-43924-V.

Molecular Weight of p-MEK-3/6: 40/37 kDa.

STORAGE

Store at 4° C, ****DO NOT FREEZE****. Stable for one year from the date of shipment. Non-hazardous. No MSDS required.

DATA



p-MEK-3/6 (B-9): sc-8407. Western blot analysis of MEK-3/6 phosphorylation in COLO 320DM (A), K-562 (B), NIH/3T3 (C), RAW 264.7 (D), KNRK (E) and L8 (F) whole cell lysates.

p-MEK-3/6 (B-9): sc-8407. Immunofluorescence staining of formalin-fixed HeLa cells showing nuclear and cytoplasmic localization (A). Immunoperoxidase staining of formalin fixed, paraffin-embedded human skeletal muscle tissue showing cytoplasmic staining of myocytes (B).

SELECT PRODUCT CITATIONS

- Wang, D., et al. 2001. Nuclear factor κ B activation by the CXC chemokine melanoma growth-stimulatory activity/growth-regulated protein involves the MEK1/p38 mitogen-activated protein kinase pathway. *J. Biol. Chem.* 276: 3650-3659.
- Wu, K., et al. 2002. Flavopiridol and trastuzumab synergistically inhibit proliferation of breast cancer cells: association with selective cooperative inhibition of cyclin D1-dependent kinase and Akt signaling pathways. *Mol. Cancer Ther.* 1: 695-706.
- Roos-Mattjus, P., et al. 2003. Phosphorylation of human Rad9 is required for genotoxin-activated checkpoint signaling. *J. Biol. Chem.* 278: 24428-24437.
- Falsig, J., et al. 2004. Specific modulation of astrocyte inflammation by inhibition of mixed lineage kinases with CEP-1347. *J. Immunol.* 173: 2762-2770.
- Alvarez, M.E., et al. 2006. Neutrophil signaling pathways activated by bacterial DNA stimulation. *J. Immunol.* 177: 4037-4046.
- Prickett, T.D., et al. 2007. Cytokine activation of p38 mitogen-activated protein kinase and apoptosis is opposed by α -4 targeting of protein phosphatase 2A for site-specific dephosphorylation of MEK-3. *Mol. Cell. Biol.* 27: 4217-4227.
- Xia, Z.P., et al. 2009. Direct activation of protein kinases by unanchored polyubiquitin chains. *Nature* 461: 114-119.
- Banh, S. and Hales, B.F. 2013. Hydroxyurea exposure triggers tissue-specific activation of p38 mitogen-activated protein kinase signaling and the DNA damage response in organogenesis-stage mouse embryos. *Toxicol. Sci.* 133: 298-308.
- Saha, K., et al. 2014. p38 δ regulates p53 to control p21^{Cip1} expression in human epidermal keratinocytes. *J. Biol. Chem.* 289: 11443-11453.

RESEARCH USE

For research use only, not for use in diagnostic procedures.