



## ZNF496 siRNA (h): sc-88120

### BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF496 (zinc finger protein 496), also known as ZKSCAN17 or NIZP1, is a 587 amino acid member of the Krüppel C<sub>2</sub>H<sub>2</sub>-type zinc-finger protein family and is thought to act as a transcriptional repressor. Localized to the nucleus, ZNF496 contains one SCAN box domain, one KRAB domain and five C<sub>2</sub>H<sub>2</sub>-type zinc fingers through which it may convey DNA, RNA and protein binding capabilities.

### REFERENCES

- Thiesen, H.J. 1990. Multiple genes encoding zinc finger domains are expressed in human T cells. *New Biol.* 2: 363-374.
- Rousseau-Merck, M.F., Hillion, J., Jonveaux, P., Couillin, P., Seite, P., Thiesen, H.J. and Berger, R. 1994. Chromosomal localization of 9 KOX zinc finger genes: physical linkages suggest clustering of KOX genes on chromosomes 12, 16, and 19. *Hum. Genet.* 92: 583-587.
- Williams, A.J., Blacklow, S.C. and Collins, T. 1999. The zinc finger-associated SCAN box is a conserved oligomerization domain. *Mol. Cell. Biol.* 19: 8526-8535.
- Rousseau-Merck, M.F., Koczan, D., Legrand, I., Möller, S., Autran, S. and Thiesen, H.J. 2003. The KOX zinc finger genes: genome wide mapping of 368 ZNF PAC clones with zinc finger gene clusters predominantly in 23 chromosomal loci are confirmed by human sequences annotated in Ensembl. *Cytogenet. Genome Res.* 98: 147-153.
- Englbrecht, C.C., Schoof, H. and Böhm, S. 2004. Conservation, diversification and expansion of C<sub>2</sub>H<sub>2</sub> zinc finger proteins in the *Arabidopsis thaliana* genome. *BMC Genomics* 5: 39-39.
- Nielsen, A.L., Jørgensen, P., Lerouge, T., Cerviño, M., Chambon, P. and Losson, R. 2004. Nizp1, a novel multitype zinc finger protein that interacts with the NSD1 histone lysine methyltransferase through a unique C2HR motif. *Mol. Cell. Biol.* 24: 5184-5196.

### CHROMOSOMAL LOCATION

Genetic locus: ZNF496 (human) mapping to 1q44.

### PRODUCT

ZNF496 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZNF496 shRNA Plasmid (h): sc-88120-SH and ZNF496 shRNA (h) Lentiviral Particles: sc-88120-V as alternate gene silencing products.

For independent verification of ZNF496 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-88120A, sc-88120B and sc-88120C.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

### APPLICATIONS

ZNF496 siRNA (h) is recommended for the inhibition of ZNF496 expression in human cells.

### SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

### RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZNF496 gene expression knockdown using RT-PCR Primer: ZNF496 (h)-PR: sc-88120-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.