



HHLA3 siRNA (h): sc-88191

BACKGROUND

Human endogenous retroviruses (HERVs) are repetitive elements that are derived from ancient germline retroviral infections. Due to their ability to move and insert next to certain genes and alter expression patterns, HERVs have been linked to several chronic diseases such as nervous systemic diseases, cancer, autoimmune and connective tissue diseases. The HERV-H family is the most abundant HERV family and has been implicated in the expression of a variety of adjacent genes. Proteins belonging to the HERV-H family are divided into one major and two minor groups based on sequence divergence. As a member of the HERV-H family, HHLA3 (HERV-H LTR-associating protein 3) is a 153 amino acid protein that is primarily expressed in liver and kidney. There are three isoforms of HHLA3 that are produced as a result of alternative splicing events.

REFERENCES

1. Feuchter-Murthy, A.E., Freeman, J.D. and Mager, D.L. 1993. Splicing of a human endogenous retrovirus to a novel phospholipase A2 related gene. *Nucleic Acids Res.* 21: 135-143.
2. Urnovitz, H.B. and Murphy, W.H. 1996. Human endogenous retroviruses: nature, occurrence, and clinical implications in human disease. *Clin. Microbiol. Rev.* 9: 72-99.
3. Kowalski, P.E., Freeman, J.D. and Mager, D.L. 1999. Intergenic splicing between a HERV-H endogenous retrovirus and two adjacent human genes. *Genomics* 57: 371-379.
4. Mager, D.L., Hunter, D.G., Schertzer, M. and Freeman, J.D. 1999. Endogenous retroviruses provide the primary polyadenylation signal for two new human genes (HHLA2 and HHLA3). *Genomics* 59: 255-263.
5. Online Mendelian Inheritance in Man, OMIM™. 1999. Johns Hopkins University, Baltimore, MD. MIM Number: 604371. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Jern, P., Lindeskog, M., Karlsson, D. and Blomberg, J. 2002. Full-length HERV-H elements with env SU open reading frames in the human genome. *AIDS Res. Hum. Retroviruses* 18: 671-676.
7. Yi, J.M. and Kim, H.S. 2004. Evolutionary implication of human endogenous retrovirus HERV-H family. *J. Hum. Genet.* 49: 215-219.
8. Yi, J.M., Kim, H.M. and Kim, H.S. 2006. Human endogenous retrovirus HERV-H family in human tissues and cancer cells: expression, identification, and phylogeny. *Cancer Lett.* 231: 228-239.
9. Balada, E., Ordi-Ros, J. and Vilardell-Tarres, M. 2009. Molecular mechanisms mediated by human endogenous retroviruses (HERVs) in autoimmunity. *Rev. Med. Virol.* 19: 273-286.

CHROMOSOMAL LOCATION

Genetic locus: HHLA3 (human) mapping to 1p31.1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

HHLA3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see HHLA3 shRNA Plasmid (h): sc-88191-SH and HHLA3 shRNA (h) Lentiviral Particles: sc-88191-V as alternate gene silencing products.

For independent verification of HHLA3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-88191A, sc-88191B and sc-88191C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

HHLA3 siRNA (h) is recommended for the inhibition of HHLA3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor HHLA3 gene expression knockdown using RT-PCR Primer: HHLA3 (h)-PR: sc-88191-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.