

NYD-SP29 siRNA (h): sc-88291

BACKGROUND

NYD-SP29 (testis development protein NYD-SP29), also known as WDR63 (WD repeat-containing protein 63) is an 891 amino acid protein that exists as two isoforms produced by alternative splicing events. NYD-SP29 contains five WD repeats and is expressed in bronchial epithelium and testis. The gene encoding NYD-SP29 maps to human chromosome 1p22.3 and mouse chromosome 3 H2. Human chromosome 1 is the largest human chromosome, spanning about 260 million base pairs and making up 8% of the human genome. There are about 3,000 genes on chromosome 1, including a large number of disease-associated genes such as those involved in familial adenomatous polyposis, Stickler syndrome, Parkinson's disease, Gaucher disease, schizophrenia and Usher syndrome. Aberrations in chromosome 1 are found in a variety of cancers, including head and neck cancer, malignant melanoma and multiple myeloma.

REFERENCES

1. Watson, M.L., Kingsmore, S.F., Johnston, G.I., Siegelman, M.H., Le Beau, M.M., Lemons, R.S., Bora, N.S., Howard, T.A., Weissman, I.L. and McEver, R.P. 1990. Genomic organization of the selectin family of leukocyte adhesion molecules on human and mouse chromosome 1. *J. Exp. Med.* 172: 263-272.
2. Blackwood, D.H., Fordyce, A., Walker, M.T., St Clair, D.M., Porteous, D.J. and Muir, W.J. 2001. Schizophrenia and affective disorders—cosegregation with a translocation at chromosome 1q42 that directly disrupts brain-expressed genes: clinical and P300 findings in a family. *Am. J. Hum. Genet.* 69: 428-433.
3. Weise, A., Starke, H., Mrasek, K., Claussen, U. and Liehr, T. 2005. New insights into the evolution of chromosome 1. *Cytogenet. Genome Res.* 108: 217-222.
4. Marzin, Y., Jamet, D., Douet-Guilbert, N., Morel, F., Le Bris, M.J., Morice, P., Abgrall, J.F., Berthou, C. and De Braekeleer, M. 2006. Chromosome 1 abnormalities in multiple myeloma. *Anticancer Res.* 26: 953-959.
5. Loneragan, K.M., Chari, R., Deleeuw, R.J., Shadoo, A., Chi, B., Tsao, M.S., Jones, S., Marra, M., Ling, V., Ng, R., Macaulay, C., Lam, S. and Lam, W.L. 2006. Identification of novel lung genes in bronchial epithelium by serial analysis of gene expression. *Am. J. Respir. Cell Mol. Biol.* 35: 651-661.
6. Zariwala, M.A., Knowles, M.R. and Omeran, H. 2007. Genetic defects in ciliary structure and function. *Annu. Rev. Physiol.* 69: 423-450.

CHROMOSOMAL LOCATION

Genetic locus: WDR63 (human) mapping to 1p22.3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

NYD-SP29 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NYD-SP29 shRNA Plasmid (h): sc-88291-SH and NYD-SP29 shRNA (h) Lentiviral Particles: sc-88291-V as alternate gene silencing products.

For independent verification of NYD-SP29 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-88291A, sc-88291B and sc-88291C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NYD-SP29 siRNA (h) is recommended for the inhibition of NYD-SP29 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NYD-SP29 gene expression knockdown using RT-PCR Primer: NYD-SP29 (h)-PR: sc-88291-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.