



connexin 59 siRNA (h): sc-88510

BACKGROUND

The connexin family of proteins form hexameric complexes called connexons that facilitate movement of low molecular weight proteins between cells via gap junctions. Connexin proteins share a common topology of four transmembrane α -helical domains, two extracellular loops, a cytoplasmic loop and cytoplasmic N- and C-termini. Many of the key functional differences between connexins arise from specific amino acid substitutions in the most highly conserved domains: the transmembrane and extracellular regions. Connexin 59, also known as connexin 58, GJA9 (Gap junction α -9 protein) or GJA10 (Gap junction α -10 protein), is a 515 amino acid multi-pass membrane protein that belongs to the connexin family. Localized to the cell membrane and to gap junctions, connexin 59 functions as part of a connexon complex through which low molecular weight materials diffuse from one cell to another.

REFERENCES

1. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 611923. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
2. Söhl, G., et al. 2003. Expression profiles of the novel human connexin genes hCx30.2, hCx40.1, and hCx62 differ from their putative mouse orthologues. *Cell Commun. Adhes.* 10: 27-36.
3. McLachlan, E., et al. 2003. Zebrafish Cx35: cloning and characterization of a gap junction gene highly expressed in the retina. *J. Neurosci. Res.* 73: 753-764.
4. Danik, S.B., et al. 2004. Modulation of cardiac gap junction expression and arrhythmic susceptibility. *Circ. Res.* 95: 1035-1041.
5. Chadjichristos, C.E., et al. 2007. Connexins: new genes in atherosclerosis. *Ann. Med.* 39: 402-411.
6. Yeager, M., et al. 2007. Gap junction channel structure in the early 21st century: facts and fantasies. *Curr. Opin. Cell Biol.* 19: 521-528.
7. Iacobas, D.A., et al. 2007. Connexin-dependent transcellular transcriptomic networks in mouse brain. *Prog. Biophys. Mol. Biol.* 94: 169-185.
8. Cotrina, M.L., et al. 2008. Adhesive properties of connexin hemichannels. *Glia* 56: 1791-1798.

CHROMOSOMAL LOCATION

Genetic locus: GJA9 (human) mapping to 1p34.3.

PRODUCT

connexin 59 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see connexin 59 shRNA Plasmid (h): sc-88510-SH and connexin 59 shRNA (h) Lentiviral Particles: sc-88510-V as alternate gene silencing products.

For independent verification of connexin 59 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-88510A, sc-88510B and sc-88510C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

connexin 59 siRNA (h) is recommended for the inhibition of connexin 59 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor connexin 59 gene expression knockdown using RT-PCR Primer: connexin 59 (h)-PR: sc-88510-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.