

ANKRD13C siRNA (h): sc-88717

BACKGROUND

Ankyrins are membrane adaptor molecules that play important roles in coupling integral membrane proteins to the spectrin-based cytoskeleton network. Mutations of ankyrin genes can lead to severe genetic diseases, such as fatal cardiac arrhythmias and hereditary spherocytosis. ANKRD13C (ankyrin repeat domain 13C) is a 541 amino acid protein that contains three ANK repeats. Conserved in chimpanzee, canine, mouse, rat, chicken, zebrafish, fruit fly, mosquito, *Caenorhabditis elegans* and *Arabidopsis thaliana*, ANKRD13C exists as three alternatively spliced isoforms. The gene that encodes ANKRD13C maps to human chromosome 1p31.1, the largest human chromosome, which makes up approximately 8% of the human genome and contains 260 million base pairs encoding 3,000 genes. Numerous diseases are linked to chromosome 1, notably the rare aging disease Hutchinson-Gilford progeria. Stickler syndrome, Parkinson's disease, Gaucher disease and Usher syndrome are also associated with chromosome 1. Aberrations in chromosome 1 exist in a variety of cancers, including head and neck cancer, malignant melanoma and multiple myeloma.

REFERENCES

- Weise, A., Starke, H., Mrasek, K., Claussen, U. and Liehr, T. 2005. New insights into the evolution of chromosome 1. *Cytogenet. Genome Res.* 108: 217-222.
- Marzin, Y., Jamet, D., Douet-Guilbert, N., Morel, F., Le Bris, M.J., Morice, P., Abgrall, J.F., Berthou, C. and De Braekeleer, M. 2006. Chromosome 1 abnormalities in multiple myeloma. *Anticancer Res.* 26: 953-959.
- Gregory, S.G., Barlow, K.F., McLay, K.E., Kaul, R., Swarbreck, D., Dunham, A., Scott, C.E., Howe, K.L., Woodfine, K., Spencer, C.C., Jones, M.C., Gillson, C., Searle, S., Zhou, Y., Kokocinski, F., McDonald, L., et al. 2006. The DNA sequence and biological annotation of human chromosome 1. *Nature* 441: 315-321.
- McClintock, D., Gordon, L.B. and Djabali, K. 2006. Hutchinson-Gilford progeria mutant Lamin A primarily targets human vascular cells as detected by an anti-Lamin A G608G antibody. *Proc. Natl. Acad. Sci. USA* 103: 2154-2159.
- Bowden, N.A., Croft, A. and Scott, R.J. 2007. Gene expression profiling in familial adenomatous polyposis adenomas and desmoid disease. *Hered. Cancer Clin. Pract.* 5: 79-96.
- Jelinsky, S.A., Choe, S.E., Crabtree, J.S., Cotreau, M.M., Wilson, E., Saraf, K., Dörner, A.J., Brown, E.L., Peano, B.J., Zhang, X., Winneker, R.C. and Harris, H.A. 2008. Molecular analysis of the vaginal response to estrogens in the ovariectomized rat and postmenopausal woman. *BMC Med. Genomics* 1: 27.
- Kirpich, I.A., Gobejishvili, L.N., Homme, M.B., Waigel, S., Cave, M., Arteel, G., Barve, S.S., McClain, C.J. and Deaciuc, I.V. 2011. Integrated hepatic transcriptome and proteome analysis of mice with high-fat diet-induced nonalcoholic fatty liver disease. *J. Nutr. Biochem.* 22: 38-45.
- SWISS-PROT/TrEMBL (Q8N6S4). World Wide Web URL: <http://www.uniprot.org/uniprot/Q8N6S4>

CHROMOSOMAL LOCATION

Genetic locus: ANKRD13C (human) mapping to 1p31.1.

PRODUCT

ANKRD13C siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ANKRD13C shRNA Plasmid (h): sc-88717-SH and ANKRD13C shRNA (h) Lentiviral Particles: sc-88717-V as alternate gene silencing products.

For independent verification of ANKRD13C (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-88717A, sc-88717B and sc-88717C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ANKRD13C siRNA (h) is recommended for the inhibition of ANKRD13C expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ANKRD13C gene expression knockdown using RT-PCR Primer: ANKRD13C (h)-PR: sc-88717-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.