

TSSK 3 siRNA (h): sc-88779

BACKGROUND

The phosphorylation and dephosphorylation of proteins on serine and threonine residues is an essential means of regulating a broad range of cellular functions in eukaryotes, including cell division, homeostasis and apoptosis. A group of proteins that are intimately involved in this process are the serine/threonine (Ser/Thr) protein kinases. TSSK 3 (testis-specific serine/threonine-protein kinase 3), also known as SPOGA3, STK22C or STK22D, is a 268 amino acid protein that contains one protein kinase domain and belongs to the Ser/Thr protein kinase family. Using magnesium as a cofactor, TSSK 3 catalyzes the ATP-dependent phosphorylation of target proteins and is thought to be involved in male germ cell development and mature sperm function. Human TSSK 3 shares 100% homology with its mouse ortholog, implying a highly conserved function between species.

REFERENCES

1. Hanks, S.K., Quinn, A.M. and Hunter, T. 1988. The protein kinase family: conserved features and deduced phylogeny of the catalytic domains. *Science* 241: 42-52.
2. Hanks, S.K. and Quinn, A.M. 1991. Protein kinase catalytic domain sequence database: identification of conserved features of primary structure and classification of family members. *Meth. Enzymol.* 200: 38-62.
3. Kueng, P., Nikolova, Z., Djonov, V., Hemphill, A., Rohrbach, V., Boehlen, D., Zuercher, G., Andres, A.C. and Ziemiecki, A. 1997. A novel family of serine/threonine kinases participating in spermiogenesis. *J. Cell Biol.* 139: 1851-1859.
4. Zuercher, G., Rohrbach, V., Andres, A.C. and Ziemiecki, A. 2000. A novel member of the testis specific serine kinase family, tssk-3, expressed in the Leydig cells of sexually mature mice. *Mech. Dev.* 93: 175-177.
5. Visconti, P.E., Hao, Z., Purdon, M.A., Stein, P., Balsara, B.R., Testa, J.R., Herr, J.C., Moss, S.B. and Kopf, G.S. 2001. Cloning and chromosomal localization of a gene encoding a novel serine/threonine kinase belonging to the subfamily of testis-specific kinases. *Genomics* 77: 163-170.
6. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 607660. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Hao, Z., Jha, K.N., Kim, Y.H., Vemuganti, S., Westbrook, V.A., Chertihin, O., Markgraf, K., Flickinger, C.J., Coppola, M., Herr, J.C. and Visconti, P.E. 2004. Expression analysis of the human testis-specific serine/threonine kinase (TSSK) homologues. A TSSK member is present in the equatorial segment of human sperm. *Mol. Hum. Reprod.* 10: 433-444.
8. Bucko-Justyna, M., Lipinski, L., Burgering, B.M. and Trzeciak, L. 2005. Characterization of testis-specific serine-threonine kinase 3 and its activation by phosphoinositide-dependent kinase-1-dependent signalling. *FEBS J.* 272: 6310-6323.
9. Xu, B., Hao, Z., Jha, K.N., Digilio, L., Urekar, C., Kim, Y.H., Pulido, S., Flickinger, C.J. and Herr, J.C. 2007. Validation of a testis specific serine/threonine kinase [TSSK] family and the substrate of TSSK 1 & 2, TSXS, as contraceptive targets. *Soc. Reprod. Fertil. Suppl.* 63: 87-101.

CHROMOSOMAL LOCATION

Genetic locus: TSSK3 (human) mapping to 1p35.1.

PRODUCT

TSSK 3 siRNA (h) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TSSK 3 shRNA Plasmid (h): sc-88779-SH and TSSK 3 shRNA (h) Lentiviral Particles: sc-88779-V as alternate gene silencing products.

For independent verification of TSSK 3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-88779A and sc-88779B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TSSK 3 siRNA (h) is recommended for the inhibition of TSSK 3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TSSK 3 gene expression knockdown using RT-PCR Primer: TSSK 3 (h)-PR: sc-88779-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.