



# OR6K6 shRNA (h) Lentiviral Particles: sc-88796-V

## BACKGROUND

Olfactory receptors are G protein-coupled receptor proteins that localize to the cilia of olfactory sensory neurons where they display affinity for and bind to a variety of odor molecules. The genes encoding olfactory receptors comprise the largest family in the human genome. The binding of olfactory receptor proteins to odor molecules triggers a signal transduction cascade that leads to the production of cAMP via an olfactory-enriched adenylate cyclase. This event ultimately leads to transmission of action potentials to the brain and the subsequent perception of smell. OR6K6 (olfactory receptor 6K6), also known as olfactory receptor OR1-21, is a 343 amino acid multi-pass membrane protein that functions as an odorant receptor, effectively binding odor molecules and initiating the propagation of signals to the primary olfactory cortex.

## REFERENCES

1. Buck, L.B. 1993. Receptor diversity and spatial patterning in the mammalian olfactory system. *Ciba Found. Symp.* 179: 51-64.
2. Sullivan, S.L., Ressler, K.J. and Buck, L.B. 1994. Odorant receptor diversity and patterned gene expression in the mammalian olfactory epithelium. *Prog. Clin. Biol. Res.* 390: 75-84.
3. Sullivan, S.L. and Dryer, L. 1996. Information processing in mammalian olfactory system. *J. Neurobiol.* 30: 20-36.
4. Touhara, K., Sengoku, S., Inaki, K., Tsuboi, A., Hirono, J., Sato, T., Sakano, H. and Haga, T. 1999. Functional identification and reconstitution of an odorant receptor in single olfactory neurons. *Proc. Natl. Acad. Sci. USA* 96: 4040-4045.
5. Kajiya, K., Inaki, K., Tanaka, M., Haga, T., Kataoka, H. and Touhara, K. 2001. Molecular bases of odor discrimination: reconstitution of olfactory receptors that recognize overlapping sets of odorants. *J. Neurosci.* 21: 6018-6025.
6. Touhara, K. 2001. Functional cloning and reconstitution of vertebrate odorant receptors. *Life Sci.* 68: 2199-2206.
7. Touhara, K. 2002. Odor discrimination by G protein-coupled olfactory receptors. *Microsc. Res. Tech.* 58: 135-141.
8. Malnic, B., Godfrey, P.A. and Buck, L.B. 2004. The human olfactory receptor gene family. *Proc. Natl. Acad. Sci. USA* 101: 2584-2589.
9. Keller, A. and Vosshall, L.B. 2008. Better smelling through genetics: mammalian odor perception. *Curr. Opin. Neurobiol.* 18: 364-369.

## RESEARCH USE

The purchase of this product conveys to the buyer the nontransferable right to use the purchased amount of the product and all replicates and derivatives for research purposes conducted by the buyer in his laboratory only (whether the buyer is an academic or for-profit entity). The buyer cannot sell or otherwise transfer (a) this product (b) its components or (c) materials made using this product or its components to a third party, or otherwise use this product or its components or materials made using this product or its components for Commercial Purposes.

## CHROMOSOMAL LOCATION

Genetic locus: OR6K6 (human) mapping to 1q23.1.

## PRODUCT

OR6K6 shRNA (h) Lentiviral Particles is a pool of concentrated, transduction-ready viral particles containing 3 target-specific constructs that encode 19-25 nt (plus hairpin) shRNA designed to knock down gene expression. Each vial contains 200  $\mu$ l frozen stock containing  $1.0 \times 10^6$  infectious units of virus (IFU) in Dulbecco's Modified Eagle's Medium with 25 mM HEPES pH 7.3. Suitable for 10-20 transductions. Also see OR6K6 siRNA (h): sc-88796 and OR6K6 shRNA Plasmid (h): sc-88796-SH as alternate gene silencing products.

## APPLICATIONS

OR6K6 shRNA (h) Lentiviral Particles is recommended for the inhibition of OR6K6 expression in human cells.

## SUPPORT REAGENTS

Control shRNA Lentiviral Particles: sc-108080. Available as 200  $\mu$ l frozen viral stock containing  $1.0 \times 10^6$  infectious units of virus (IFU); contains an shRNA construct encoding a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor OR6K6 gene expression knockdown using RT-PCR Primer: OR6K6 (h)-PR: sc-88796-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## BIOSAFETY

Lentiviral particles can be employed in standard Biosafety Level 2 tissue culture facilities (and should be treated with the same level of caution as with any other potentially infectious reagent). Lentiviral particles are replication-incompetent and are designed to self-inactivate after transduction and integration of shRNA constructs into genomic DNA of target cells.

## STORAGE

Store lentiviral particles at -80° C. Stable for at least one year from the date of shipment. Once thawed, particles can be stored at 4° C for up to one week. Avoid repeated freeze thaw cycles.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.