SANTA CRUZ BIOTECHNOLOGY, INC.

PDCL2 siRNA (h): sc-89178



BACKGROUND

PDCL2 (phosducin-like protein 2) is a 241 amino acid protein that belongs to the phosducin family. While it forms a complex with 14-3-3 proteins, PDCL2 may play a role in germ cell maturation. As a member of the phosducin-like protein family, PDCL2 is a putative modulator of heterotrimeric G proteins. More specifically, members of the phosducin-like protein family have been shown to bind to the β - γ subunits of G proteins. The gene that encodes PDCL2 consists of more than 35,000 bases and maps to human chromosome 4q12. Chromosome 4 represents approximately 6% of the human genome and contains nearly 900 genes. Notably, the Huntingtin gene, which is found to encode an expanded glutamine tract in cases of Huntington's disease, is encoded by a gene that maps to chromosome 4. FGFR-3 is also encoded by a gene located on chromosome 4 and has been associated with thanatophoric dwarfism, achondroplasia, Muenke syndrome and bladder cancer. Chromosome 4 is also tied to Ellis-van Creveld syndrome, methylmalonic acidemia and polycystic kidney disease.

REFERENCES

- Bonaventure, J., et al. 1996. Common mutations in the fibroblast growth factor receptor 3 (FGFR 3) gene account for achondroplasia, hypochondroplasia, and thanatophoric dwarfism. Am. J. Med. Genet. 63: 148-154.
- 2. Kalchman, M.A., et al. 1996. Huntingtin is ubiquitinated and interacts with a specific ubiquitin-conjugating enzyme. J. Biol. Chem. 271: 19385-19394.
- Singhrao, S.K., et al. 1998. Huntingtin protein colocalizes with lesions of neurodegenerative diseases: an investigation in Huntington's, Alzheimer's, and Pick's diseases. Exp. Neurol. 150: 213-222.
- Krakow, D., et al. 2000. Exclusion of the Ellis-van Creveld region on chromosome 4p16 in some families with asphyxiating thoracic dystrophy and short-rib polydactyly syndromes. Eur. J. Hum. Genet. 8: 645-648.
- Sommardahl, C., et al. 2001. Phenotypic variations of orpk mutation and chromosomal localization of modifiers influencing kidney phenotype. Physiol. Genomics 7: 127-134.
- Dobson, C.M., et al. 2002. Identification of the gene responsible for the cbIA complementation group of vitamin B12-responsive methylmalonic acidemia based on analysis of prokaryotic gene arrangements. Proc. Natl. Acad. Sci. USA 99: 15554-15559.
- 7. Lopez, P., et al. 2003. A novel germ line-specific gene of the phosducinlike protein (PhLP) family. A meiotic function conserved from yeast to mice. J. Biol. Chem. 278: 1751-1757.
- 8. Online Mendelian Inheritance in Man, OMIM™. 2007. Johns Hopkins University, Baltimore, MD. MIM Number: 611676. World Wide Web URL: http://www.ncbi.nlm.nih.gov/omim/
- Lou, X., et al. 2009. Structure of the thioredoxin-fold domain of human phosducin-like protein 2. Acta Crystallogr. Sect. F Struct. Biol. Cryst. Commun. 65: 67-70.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: PDCL2 (human) mapping to 4q12.

PRODUCT

PDCL2 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PDCL2 shRNA Plasmid (h): sc-89178-SH and PDCL2 shRNA (h) Lentiviral Particles: sc-89178-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNAse-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNAse-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

PDCL2 siRNA (h) is recommended for the inhibition of PDCL2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PDCL2 gene expression knockdown using RT-PCR Primer: PDCL2 (h)-PR: sc-89178-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.