

TCAF1 siRNA (h): sc-89631

BACKGROUND

TRPM8 (transient receptor potential cation channel, subfamily M, member 8) is a multi-pass membrane protein that is expressed at high levels in prostate. TRPM8 functions as a receptor-activated cation channel that is permeable to monovalent cations (such as sodium and potassium) and divalent Ca^{2+} and is involved in the detection of temperature sensations (such as the feeling of coolness) throughout the body. TRPM8 is overexpressed in prostate tumors, as well as in colon, breast and lung cancers. TCAF1 (TRPM8 channel-associated factor 1), also known as FAM115A, is a 921 amino acid plasma membrane protein that is expressed as two alternatively spliced isoforms and belong to the TCAF family. TCAF1 isoform 2 is expressed in prostate and strongly expressed in cancerous prostate tissue. TCAF1 consists of a peptidase M60 domain and a C-terminal region that is necessary for the channel activity stimulation. TCAF1 along with family member TCAF2 interact with plasma membrane cation channel TRPM8 and promotes its trafficking to the cell surface. TCAF1 is encoded by a gene located on human chromosome 7q35.

REFERENCES

1. Tsavaler, L., Shaper, M.H., Morkowski, S. and Laus, R. 2001. TRPP8, a novel prostate-specific gene, is upregulated in prostate cancer and other malignancies and shares high homology with transient receptor potential calcium channel proteins. *Cancer Res.* 61: 3760-3769.
2. de la Peña, E., Mäklä, A., Cabedo, H., Belmonte, C. and Viana, F. 2005. The contribution of TRPM8 channels to cold sensing in mammalian neurons. *J. Physiol.* 567: 415-426.
3. Erler, I., Al-Ansary, D.M., Wissenbach, U., Wagner, T.F., Flockerzi, V. and Niemeyer, B.A. 2006. Trafficking and assembly of the cold-sensitive TRPM8 channel. *J. Biol. Chem.* 281: 38396-38404.
4. Facer, P., Casula, M.A., Smith, G.D., Benham, C.D., Chessell, I.P., Bountra, C., Sinisi, M., Birch, R. and Anand, P. 2007. Differential expression of the capsaicin receptor TRPV1 and related novel receptors TRPV3, TRPV4 and TRPM8 in normal human tissues and changes in traumatic and diabetic neuropathy. *BMC Neurol.* 7: 11.
5. Bödding, M., Wissenbach, U. and Flockerzi, V. 2007. Characterisation of TRPM8 as a pharmacophore receptor. *Cell Calcium.* 42: 618-628.
6. Kühn, F.J., Knop, G. and Lückhoff, A. 2007. The transmembrane segment S6 determines cation versus anion selectivity of TRPM2 and TRPM8. *J. Biol. Chem.* 282: 27598-27609.
7. Mahieu, F., Owsianik, G., Verbert, L., Janssens, A., De Smedt, H., Nilius, B. and Voets, T. 2007. TRPM8-independent menthol-induced Ca^{2+} release from endoplasmic reticulum and Golgi. *J. Biol. Chem.* 282: 3325-3336.
8. Bidaux, G., Flourakis, M., Thebault, S., Zholos, A., Beck, B., Gkika, D., Roud-baraki, M., Bonnal, J.L., Mauroy, B., Shuba, Y., Skryma, R. and Prevarskaya, N. 2007. Prostate cell differentiation status determines transient receptor potential melastatin member 8 channel subcellular localization and function. *J. Clin. Invest.* 117: 1647-1657.

CHROMOSOMAL LOCATION

Genetic locus: TCAF1 (human) mapping to 7q35.

PRODUCT

TCAF1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TCAF1 shRNA Plasmid (h): sc-89631-SH and TCAF1 shRNA (h) Lentiviral Particles: sc-89631-V as alternate gene silencing products.

For independent verification of TCAF1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-89631A, sc-89631B and sc-89631C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20°C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20°C , avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TCAF1 siRNA (h) is recommended for the inhibition of TCAF1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl . Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TCAF1 gene expression knockdown using RT-PCR Primer: TCAF1 (h)-PR: sc-89631-PR (20 μl). Annealing temperature for the primers should be $55-60^{\circ}\text{C}$ and the extension temperature should be $68-72^{\circ}\text{C}$.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.