

GLCCI1 siRNA (h): sc-89671

BACKGROUND

GLCCI1 (glucocorticoid induced transcript 1), also known as TSSN1, GIG18 or FAM117C, is a 547 amino acid protein found in thymus and CD4⁺CD8⁺ cells during specific stages of spermatogenesis in testis. Subject to phosphorylation, GLCCI1 is encoded by a gene located on human chromosome 7, which houses over 1,000 genes and comprises nearly 5% of the human genome. Chromosome 7 has been linked to osteogenesis imperfecta, Pendred syndrome, lissencephaly, citrullinemia and Shwachman-Diamond syndrome. The deletion of a portion of the q arm of chromosome 7 is associated with Williams-Beuren syndrome, a condition characterized by mild mental retardation, an unusual comfort and friendliness with strangers and an elfin appearance. Deletions of portions of the q arm of chromosome 7 are also seen in a number of myeloid disorders including cases of acute myelogenous leukemia and myelodysplasia.

REFERENCES

1. Tsipouras, P., Myers, J.C., Ramirez, F. and Prockop, D.J. 1983. Restriction fragment length polymorphism associated with the pro α 2(I) gene of human type I procollagen. Application to a family with an autosomal dominant form of Osteogenesis imperfecta. *J. Clin. Invest.* 72: 1262-1267.
2. Liang, H., Fairman, J., Claxton, D.F., Nowell, P.C., Green, E.D. and Nagarajan, L. 1998. Molecular anatomy of chromosome 7q deletions in myeloid neoplasms: evidence for multiple critical loci. *Proc. Natl. Acad. Sci. USA* 95: 3781-3785.
3. Iwasaki, S., Usami, S., Abe, S., Isoda, H., Watanabe, T. and Hoshino, T. 2001. Long-term audiological feature in Pendred syndrome caused by PDS mutation. *Arch. Otolaryngol. Head Neck Surg.* 127: 705-708.
4. Miazek, A. and Malissen, B. 2003. Two genes, three messengers: hybrid transcript between a gene expressed at specific stages of T-cell and sperm maturation and an unrelated adjacent gene. *Immunogenetics* 54: 681-692.
5. Reiner, O., Sapoznik, S. and Sapir, T. 2006. Lissencephaly 1 linking to multiple diseases: mental retardation, neurodegeneration, schizophrenia, male sterility, and more. *Neuromolecular Med.* 8: 547-565.
6. Gilbert-Dussardier, B. 2006. Williams-Beuren syndrome. *Rev. Prat.* 56: 2102-2106.
7. Leone, G., Pagano, L., Ben-Yehuda, D. and Voso, M.T. 2007. Therapy-related leukemia and myelodysplasia: susceptibility and incidence. *Haematologica* 92: 1389-1398.

CHROMOSOMAL LOCATION

Genetic locus: GLCCI1 (human) mapping to 7p21.3.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

GLCCI1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GLCCI1 shRNA Plasmid (h): sc-89671-SH and GLCCI1 shRNA (h) Lentiviral Particles: sc-89671-V as alternate gene silencing products.

For independent verification of GLCCI1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-89671A, sc-89671B and sc-89671C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GLCCI1 siRNA (h) is recommended for the inhibition of GLCCI1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GLCCI1 gene expression knockdown using RT-PCR Primer: GLCCI1 (h)-PR: sc-89671-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.