

GIMAP8 siRNA (h): sc-89799

BACKGROUND

The GTPase of the immunity-associated protein (GIMAP) family of proteins include seven members that are expressed by genes residing on human chromosome 7. GIMAP proteins have been implicated in the regulation of lymphomyeloid cell survival. GIMAP8, also known as IAN9 (immune-associated nucleotide-binding protein 9) or IANT, is a 665 amino acid protein that localizes to Golgi apparatus, Endoplasmic reticulum and mitochondria. Suggested to have an anti-apoptotic effect on the immune system, GIMAP8 plays a role in infection response and is encoded by a gene that maps to human chromosome 7q36.1. Chromosome 7 houses over 1,000 genes, comprises nearly 5% of the human genome and has been linked to Osteogenesis imperfecta, Pendred syndrome, Lissencephaly, Citrullinemia and Shwachman-Diamond syndrome.

REFERENCES

1. Tsiouras, P., Myers, J.C., Ramirez, F. and Prockop, D.J. 1983. Restriction fragment length polymorphism associated with the pro α 2(I) gene of human type I procollagen. Application to a family with an autosomal dominant form of Osteogenesis imperfecta. *J. Clin. Invest.* 72: 1262-1267.
2. Iwasaki, S., Usami, S., Abe, S., Isoda, H., Watanabe, T. and Hoshino, T. 2001. Long-term audiological feature in Pendred syndrome caused by PDS mutation. *Arch. Otolaryngol. Head Neck Surg.* 127: 705-708.
3. Cambot, M., Aresta, S., Kahn-Perlès, B., de Gunzburg, J. and Romeo, P.H. 2002. Human immune associated nucleotide 1: a member of a new guanosine triphosphatase family expressed in resting T and B cells. *Blood* 99: 3293-3301.
4. Stamm, O., Krücken, J., Schmitt-Wrede, H.P., Benten, W.P. and Wunderlich, F. 2002. Human ortholog to mouse gene *imap38* encoding an ER-localizable G protein belongs to a gene family clustered on chromosome 7q32-36. *Gene* 282: 159-167.
5. Krücken, J., Schroetel, R.M., Müller, I.U., Saïdani, N., Marinovski, P., Benten, W.P., Stamm, O. and Wunderlich, F. 2004. Comparative analysis of the human *gimap* gene cluster encoding a novel GTPase family. *Gene* 341: 291-304.
6. Dion, C., Carter, C., Hepburn, L., Coadwell, W.J., Morgan, G., Graham, M., Pugh, N., Anderson, G., Butcher, G.W. and Miller, J.R. 2005. Expression of the *Ian* family of putative GTPases during T cell development and description of an *Ian* with three sets of GTP/GDP-binding motifs. *Int. Immunol.* 17: 1257-1268.
7. Reiner, O., Sapoznik, S. and Sapir, T. 2006. Lissencephaly 1 linking to multiple diseases: mental retardation, neurodegeneration, schizophrenia, male sterility, and more. *Neuromolecular Med.* 8: 547-565.

CHROMOSOMAL LOCATION

Genetic locus: GIMAP8 (human) mapping to 7q36.1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

GIMAP8 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GIMAP8 shRNA Plasmid (h): sc-89799-SH and GIMAP8 shRNA (h) Lentiviral Particles: sc-89799-V as alternate gene silencing products.

For independent verification of GIMAP8 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-89799A, sc-89799B and sc-89799C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GIMAP8 siRNA (h) is recommended for the inhibition of GIMAP8 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GIMAP8 gene expression knockdown using RT-PCR Primer: GIMAP8 (h)-PR: sc-89799-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.