

NSUN5C siRNA (h): sc-89874

BACKGROUND

NSUN5 (NOL1/NOP2/Sun domain family, member 5), also known as NOL1, p120 or NOL1R, is a 429 amino acid protein that belongs to the methyltransferase superfamily and exists as multiple alternatively spliced isoforms. Expressed ubiquitously with higher expression in heart, placenta and skeletal muscle, NSUN5 is thought to function as an S-adenosyl-L-methionine-dependent methyl-transferase whose absence may be associated with the pathogenesis of Williams syndrome. NSUN5 is subject to post-translational phosphorylation, probably by ATM or ATR. NSUN5B and NSUN5C are truncated proteins that are also ubiquitously expressed, but with some tissue specific patterns.

REFERENCES

- Liang, H., Fairman, J., Claxton, D.F., Nowell, P.C., Green, E.D. and Nagarajan, L. 1998. Molecular anatomy of chromosome 7q deletions in myeloid neoplasms: evidence for multiple critical loci. *Proc. Natl. Acad. Sci. USA* 95: 3781-3785.
- Francke, U. 1999. Williams-Beuren syndrome: genes and mechanisms. *Hum. Mol. Genet.* 8: 1947-1954.
- Doll, A. and Grzeschik, K.H. 2001. Characterization of two novel genes, WBSCR20 and WBSCR22, deleted in Williams-Beuren syndrome. *Cytogenet. Cell Genet.* 95: 20-27.
- Merla, G., Ucla, C., Guipponi, M. and Reymond, A. 2002. Identification of additional transcripts in the Williams-Beuren syndrome critical region. *Hum. Genet.* 110: 429-438.
- Bayes, M., Magano, L.F., Rivera, N., Flores, R. and Perez Jurado, L.A. 2003. Mutational mechanisms of Williams-Beuren syndrome deletions. *Am. J. Hum. Genet.* 73: 131-151.
- Eckert, M.A., Galaburda, A.M., Mills, D.L., Bellugi, U., Korenberg, J.R. and Reiss, A.L. 2006. The neurobiology of Williams syndrome: cascading influences of visual system impairment? *Cell. Mol. Life Sci.* 63: 1867-1875.
- Osborne, L.R., Joseph-George, A.M. and Scherer, S.W. 2006. Williams-Beuren syndrome diagnosis using fluorescence *in situ* hybridization. *Methods Mol. Med.* 126:113-128.
- Schubert, C. 2009. The genomic basis of the Williams-Beuren syndrome. *Cell. Mol. Life Sci.* 66: 1178-1197.

CHROMOSOMAL LOCATION

Genetic locus: NSUN5P2 (human) mapping to 7q11.23.

PRODUCT

NSUN5C siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NSUN5C shRNA Plasmid (h): sc-89874-SH and NSUN5C shRNA (h) Lentiviral Particles: sc-89874-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NSUN5C siRNA (h) is recommended for the inhibition of NSUN5C expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

NSUN5 (H-10): sc-376147 is recommended as a control antibody for monitoring of NSUN5C gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker[™] Molecular Weight Standards: sc-2035, UltraCruz[®] Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz[®] Mounting Medium: sc-24941 or UltraCruz[®] Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NSUN5C gene expression knockdown using RT-PCR Primer: NSUN5C (h)-PR: sc-89874-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.