

GOLGA8K siRNA (h): sc-90160

BACKGROUND

The Golgi complex plays an essential role in the post-translational modification and sorting of proteins transported from the endoplasmic reticulum (ER). The Golgi stack consists of a distinct *cis* face, or entry face, and a *trans* face, or exit face, which are connected via the *cis*, medial and *trans* Golgi networks. GOLGA8K (golgin A8 family, member K) is a 607 amino acid protein belonging to the GOLGA8 family. The gene encoding GOLGA8K maps to human chromosome 15q13.3. Encoding more than 700 genes, chromosome 15 is made up of approximately 106 million base pairs and is about 3% of the human genome. Angelman and Prader-Willi syndromes are associated with loss of function or deletion of genes in the 15q11-q13 region. Tay-Sachs disease is a lethal disorder associated with mutations of the HEXA gene, which is encoded by chromosome 15. Marfan syndrome is associated with chromosome 15 through the FBN1 gene.

REFERENCES

- Hurowitz, G.I., Silver, J.M., Brin, M.F., Williams, D.T. and Johnson, W.G. 1993. Neuropsychiatric aspects of adult-onset Tay-Sachs disease: two case reports with several new findings. *J. Neuropsychiatry Clin. Neurosci.* 5: 30-36.
- Ríos, R.M., Sanchís, A., Tassin, A.M., Fedriani, C. and Bornens, M. 2004. GMAP-210 recruits γ -Tubulin complexes to *cis*-Golgi membranes and is required for Golgi ribbon formation. *Cell* 118: 323-335.
- Ouchi, M., West, K., Crabb, J.W., Kinoshita, S. and Kamei, M. 2005. Proteomic analysis of vitreous from diabetic macular edema. *Exp. Eye Res.* 81: 176-182.
- Barr, F.A. and Egerer, J. 2005. Golgi positioning: are we looking at the right MAP? *J. Cell Biol.* 168: 993-998.
- Jiang, Y.H., Wauki, K., Liu, Q., Bressler, J., Pan, Y., Kashork, C.D., Shaffer, L.G. and Beaudet, A.L. 2008. Genomic analysis of the chromosome 15q11-q13 Prader-Willi syndrome region and characterization of transcripts for GOLGA8E and WHCD1L1 from the proximal breakpoint region. *BMC Genomics* 9: 50.
- Midla, G.S. 2008. Diagnosis and management of patients with Marfan syndrome. *JAAPA* 21: 21-25.
- Dan, B. 2009. Angelman syndrome: current understanding and research prospects. *Epilepsia* 50: 2331-2339.
- Ferrer-Bolufer, I., Dalmau, J., Quiroga, R., Oltra, S., Orellana, C., Monfort, S., Roselló, M., De La Osa, A. and Martinez, F. 2009. Tyrosinemia type 1 and Angelman syndrome due to paternal uniparental isodisomy 15. *J. Inher. Metab. Dis.* 32: S349-S353.
- Wawrzik, M., Unmehopa, U.A., Swaab, D.F., van de Nes, J., Buiting, K. and Horsthemke, B. 2010. The C15orf2 gene in the Prader-Willi syndrome region is subject to genomic imprinting and positive selection. *Neurogenetics* 11: 153-161.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: GOLGA8K (human) mapping to 15q13.3.

PRODUCT

GOLGA8K siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GOLGA8K shRNA Plasmid (h): sc-90160-SH and GOLGA8K shRNA (h) Lentiviral Particles: sc-90160-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GOLGA8K siRNA (h) is recommended for the inhibition of GOLGA8K expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GOLGA8K gene expression knockdown using RT-PCR Primer: GOLGA8K (h)-PR: sc-90160-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.