

protein 4.2 siRNA (h): sc-90233

BACKGROUND

Protein 4.2, also known as Erythrocyte membrane protein band 4.2, is a 691 amino acid transmembrane protein, which possibly regulates mechanical and morphological properties of erythrocytes. Protein 4.2 functions to strongly influence CD47 levels and also regulate the association between Ankyrin and protein 3. Appearing on erythroblasts at a very late stage of red blood cell development, protein 4.2 is predominantly found in liver and is also expressed in spleen, stomach, testis and eye. One of several members of the transglutaminase family, protein 4.2 is subject to a loss of function caused by an amino acid substitution from Cys to Ala in the active site. Complete or partial protein 4.2 absence leads to a weaker than usual association of ankyrin to the membrane skeleton. Defects in the gene encoding protein 4.2 are the cause of hereditary spherocytosis, a hematologic disorder characterized by abnormally shaped erythrocytes and chronic hemolytic anemia. A short and long isoform of protein 4.2 exist as a result of an alternative splicing events. The short isoform is characterized as the major protein 4.2 species in human erythrocyte membranes.

REFERENCES

1. Sung, L.A., Chien, S., Fan, Y.S., Lin, C.C., Lambert, K., Zhu, L., Lam, J.S. and Chang, L.S. 1992. Human erythrocyte protein 4.2: isoform expression, differential splicing, and chromosomal assignment. *Blood* 79: 2763-2770.
2. Zhu, L., Kahwash, S.B. and Chang, L.S. 1998. Developmental expression of mouse erythrocyte protein 4.2 mRNA: evidence for specific expression in erythroid cells. *Blood* 91: 695-705.
3. Mouro-Chanteloup, I., Delaunay, J., Gane, P., Nicolas, V., Johansen, M., Brown, E.J., Peters, L.L., Van Kim, C.L., Cartron, J.P. and Colin, Y. 2003. Evidence that the red cell skeleton protein 4.2 interacts with the Rh membrane complex member CD47. *Blood* 101: 338-344.
4. Dahl, K.N., Parthasarathy, R., Westhoff, C.M., Layton, D.M. and Discher, D.E. 2004. Protein 4.2 is critical to CD47-membrane skeleton attachment in human red cells. *Blood* 103: 1131-1136.
5. Toye, A.M., Ghosh, S., Young, M.T., Jones, G.K., Sessions, R.B., Ramauge, M., Leclerc, P., Basu, J., Delaunay, J. and Tanner, M.J. 2005. Protein-4.2 association with band 3 (AE1, SLCA4) in *Xenopus* oocytes: effects of three natural protein-4.2 mutations associated with hemolytic anemia. *Blood* 105: 4088-4095.
6. Remus, R., Kanzaki, A., Yawata, A., Wada, H., Nakanishi, H., Sugihara, T., Zeschngk, M., Zuther, I., Schmitz, B., Naumann, F., Doerfler, W. and Yawata, Y. 2005. Relationships between DNA methylation and expression in erythrocyte membrane protein (band 3, protein 4.2, and β -spectrin) genes during human erythroid development and differentiation. *Int. J. Hematol.* 82: 422-429.
7. Su, Y., Ding, Y., Jiang, M., Jiang, W., Hu, X. and Zhang, Z. 2006. Associations of protein 4.2 with band 3 and ankyrin. *Mol. Cell. Biochem.* 289: 159-166.
8. Su, Y., Ding, Y., Jiang, M., Hu, X. and Zhang, Z. 2007. Protein 4.2 Komatsu (D175Y) associated with the lack of interaction with ankyrin in human red blood cells. *Blood Cells Mol. Dis.* 38: 221-228.

CHROMOSOMAL LOCATION

Genetic locus: EPB42 (human) mapping to 15q15.2.

PRODUCT

protein 4.2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see protein 4.2 shRNA Plasmid (h): sc-90233-SH and protein 4.2 shRNA (h) Lentiviral Particles: sc-90233-V as alternate gene silencing products.

For independent verification of protein 4.2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-90233A, sc-90233B and sc-90233C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

protein 4.2 siRNA (h) is recommended for the inhibition of protein 4.2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor protein 4.2 gene expression knockdown using RT-PCR Primer: protein 4.2 (h)-PR: sc-90233-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.