

TMCO5 siRNA (h): sc-90270

BACKGROUND

TMCO5 (transmembrane and coiled-coil domain-containing protein 5), also known as TMC05A, is a 288 amino acid single-pass membrane protein that belongs to the TMC05 family and exists as two alternatively spliced isoforms. The gene that encodes TMC05 consists of about 16,284 bases and maps to human chromosome 15q14. Housing approximately 106 million base pairs and encoding more than 700 genes, chromosome 15 makes up about 3% of the human genome. Angelman and Prader-Willi syndromes are associated with loss of function or deletion of genes in the 15q11-q13 region. In the case of Angelman syndrome, this loss is due to inactivity of the maternal 15q11-q13 encoded UBE3A gene in the brain by either chromosomal deletion or mutation. In cases of Prader-Willi syndrome, there is a partial or complete deletion of this region from the paternal copy of chromosome 15. Tay-Sachs disease is a lethal disorder associated with mutations of the HEXA gene, which is encoded by chromosome 15. Marfan syndrome is associated with chromosome 15 through the FBN1 gene.

REFERENCES

- Hurowitz, G.I., Silver, J.M., Brin, M.F., Williams, D.T. and Johnson, W.G. 1993. Neuropsychiatric aspects of adult-onset Tay-Sachs disease: two case reports with several new findings. *J. Neuropsychiatry Clin. Neurosci.* 5: 30-36.
- Midla, G.S. 2008. Diagnosis and management of patients with Marfan syndrome. *JAAPA* 21: 21-25.
- Dan, B. 2009. Angelman syndrome: current understanding and research prospects. *Epilepsia* 50: 2331-2339.
- Ferrer-Bolufer, I., Dalmau, J., Quiroga, R., Oltra, S., Orellana, C., Monfort, S., Roselló, M., De La Osa, A. and Martinez, F. 2009. Tyrosinemia type 1 and Angelman syndrome due to paternal uniparental isodisomy 15. *J. Inherit. Metab. Dis.* 32: S349-S353.
- Lanktree, M.B., Hegele, R.A., Yusuf, S. and Anand, S.S. 2009. Multi-ethnic genetic association study of carotid intima-media thickness using a targeted cardiovascular SNP microarray. *Stroke* 40: 3173-3179.
- Wawrzik, M., Unmehopa, U.A., Swaab, D.F., van de Nes, J., Buiting, K. and Horsthemke, B. 2010. The C15orf2 gene in the Prader-Willi syndrome region is subject to genomic imprinting and positive selection. *Neurogenetics* 11: 153-161.
- Bauerschlag, D.O., Ammerpohl, O., Bräutigam, K., Schem, C., Lin, Q., Weigel, M.T., Hilpert, F., Arnold, N., Maass, N., Meinhold-Heerlein, I. and Wagner, W. 2011. Progression-free survival in ovarian cancer is reflected in epigenetic DNA methylation profiles. *Oncology* 80: 12-20.

CHROMOSOMAL LOCATION

Genetic locus: TMC05A (human) mapping to 15q14.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

TMCO5 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TMC05 shRNA Plasmid (h): sc-90270-SH and TMC05 shRNA (h) Lentiviral Particles: sc-90270-V as alternate gene silencing products.

For independent verification of TMC05 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-90270A, sc-90270B and sc-90270C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TMCO5 siRNA (h) is recommended for the inhibition of TMC05 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TMC05 gene expression knockdown using RT-PCR Primer: TMC05 (h)-PR: sc-90270-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.