



# PCGF5 siRNA (h): sc-90356

## BACKGROUND

Polycomb group (PcG) proteins form multiprotein complexes that regulate expression patterns of developmental and cell proliferation genes. Several members of the PcG contain RING finger domains and are identified as a subclass of RING finger proteins. The RING-type zinc finger motif is present in a number of viral and eukaryotic proteins and is made of a conserved cysteine-rich domain that is able to bind two zinc atoms. Proteins that contain the RING-type zinc finger conserved domain are generally involved in the ubiquitination pathway of protein degradation. PCGF5 (Polycomb group RING finger protein 5), also known as RNF159 (RING finger protein 159), is a 256 amino acid transcriptional regulator that is encoded by a gene located on chromosome 10q23.32, which contains a plethora of interesting genes and represents between 4 and 4.5 percent of the total DNA in cells. Jackson-Weiss, Cowden and Usher syndromes are a few diseases related to genes on chromosome 10. Two isoforms of PCGF5 are expressed due to alternative splicing.

## REFERENCES

1. Kanno, M., Hasegawa, M., Ishida, A., Isono, K. and Taniguchi, M. 1995. Mel-18, a Polycomb group-related mammalian gene, encodes a transcriptional negative regulator with tumor suppressive activity. *EMBO J.* 14: 5672-5678.
2. Satijn, D.P., Gunster, M.J., van der Vlag, J., Hamer, K.M., Schul, W., Alkema, M.J., Saurin, A.J., Freemont, P.S., van Driel, R. and Otte, A.P. 1997. RING1 is associated with the Polycomb group protein complex and acts as a transcriptional repressor. *Mol. Cell. Biol.* 17: 4105-4113.
3. Joazeiro, C.A. and Weissman, A.M. 2000. RING finger proteins: mediators of ubiquitin ligase activity. *Cell* 102: 549-552.
4. Tuckfield, A., Clouston, D.R., Wilanowski, T.M., Zhao, L.L., Cunningham, J.M. and Jane, S.M. 2002. Binding of the RING Polycomb proteins to specific target genes in complex with the grainyhead-like family of developmental transcription factors. *Mol. Cell. Biol.* 22: 1936-1946.
5. Moore, R. and Boyd, L. 2004. Analysis of RING finger genes required for embryogenesis in *C. elegans*. *Genesis* 38: 1-12.
6. Yang, Y., Lorick, K.L., Jensen, J.P. and Weissman, A.M. 2005. Expression and evaluation of RING finger proteins. *Methods Enzymol.* 398: 103-112.

## CHROMOSOMAL LOCATION

Genetic locus: PCGF5 (human) mapping to 10q23.32.

## PRODUCT

PCGF5 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see PCGF5 shRNA Plasmid (h): sc-90356-SH and PCGF5 shRNA (h) Lentiviral Particles: sc-90356-V as alternate gene silencing products.

For independent verification of PCGF5 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-90356A, sc-90356B and sc-90356C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

PCGF5 siRNA (h) is recommended for the inhibition of PCGF5 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor PCGF5 gene expression knockdown using RT-PCR Primer: PCGF5 (h)-PR: sc-90356-PR (20  $\mu$ l, 638 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.