

ZNF365 siRNA (h): sc-90549

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF365 (zinc finger protein 365) is a 407 amino acid protein with its expression restricted to brain, lung, liver, placenta, kidney and pancreas. Overexpression of ZNF365 causes abnormal mitosis and mutant ZNF365 lacking a C-terminus disrupts γ -Tubulin localization to the nucleus. Alternative splicing results in at least four different isoforms of ZNF365, designated ZNF365A-D. A mutation in the gene encoding ZNF365 disrupts the expression of ZNF365D (also known as Talanin) and is involved in susceptibility to uric acid nephrolithiasis, a multifactorial urinary tract stone disease that is influenced by genetics and environmental factors.

REFERENCES

1. Payre, F. and Vincent, A. 1988. Finger proteins and DNA-specific recognition: distinct patterns of conserved amino acids suggest different evolutionary modes. *FEBS Lett.* 234: 245-250.
2. Thiesen, H.J. 1990. Multiple genes encoding zinc finger domains are expressed in human T cells. *New Biol.* 2: 363-374.
3. Rosenfeld, R. and Margalit, H. 1993. Zinc fingers: conserved properties that can distinguish between spurious and actual DNA-binding motifs. *J. Biomol. Struct. Dyn.* 11: 557-570.
4. Ombra, M.N., Forabosco, P., Casula, S., Angius, A., Maestrale, G., Petretto, E., Casu, G., Colussi, G., Usai, E., Melis, P. and Pirastu, M. 2001. Identification of a new candidate locus for uric acid nephrolithiasis. *Am. J. Hum. Genet.* 68: 1119-1129.
5. Online Mendelian Inheritance in Man, OMIM[™]. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 605990. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Gianfrancesco, F., Esposito, T., Ombra, M.N., Forabosco, P., Maninchedda, G., Fattorini, M., Casula, S., Vaccargiu, S., Casu, G., Cardia, F., Deiana, I., Melis, P., Falchi, M. and Pirastu, M. 2003. Identification of a novel gene and a common variant associated with uric acid nephrolithiasis in a Sardinian genetic isolate. *Am. J. Hum. Genet.* 72: 1479-1491.
7. Gianfrancesco, F., Esposito, T., Casu, G., Maninchedda, G., Roberto, R. and Pirastu, M. 2004. Emergence of Talanin protein associated with human uric acid nephrolithiasis in the Hominidae lineage. *Gene* 339: 131-138.
8. Gianfrancesco, F. and Esposito, T. 2005. Multifactorial disorder: molecular and evolutionary insights of uric acid nephrolithiasis. *Minerva Med.* 96: 409-416.

CHROMOSOMAL LOCATION

Genetic locus: ZNF365 (human) mapping to 10q21.2.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

ZNF365 siRNA (h) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZNF365 shRNA Plasmid (h): sc-90549-SH and ZNF365 shRNA (h) Lentiviral Particles: sc-90549-V as alternate gene silencing products.

For independent verification of ZNF365 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-90549A and sc-90549B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ZNF365 siRNA (h) is recommended for the inhibition of ZNF365 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZNF365 gene expression knockdown using RT-PCR Primer: ZNF365 (h)-PR: sc-90549-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.