

hnRNP H3 siRNA (h): sc-90762

BACKGROUND

Heterogeneous nuclear ribonucleoproteins (hnRNPs) constitute a set of polypeptides that contribute to mRNA transcription and pre-mRNA processing, as well as mature mRNA transport to the cytoplasm and translation. hnRNPs also bind heterogeneous nuclear RNA (hnRNA), which are the transcripts produced by RNA polymerase II. There are approximately 20 known hnRNP proteins, and their complexes are the major constituents of the spliceosome. The majority of hnRNP proteins are localized to the nucleus; however some shuttle between the nucleus and the cytoplasm. hnRNPs F and H are highly related factors that preferentially associate with poly(rG) regions on RNA. hnRNP H3, also known as hnRNP 2H9, is a 346 amino acid protein involved in RNA processing, as well as early heat shock-inducing splicing arrest. hnRNP H3 contains two RNA recognition motif (RRM) domains, which include locations for binding single-stranded RNA. hnRNP H3 is expressed as six isoforms generated by alternative splicing of the pre-mRNA.

REFERENCES

1. Mahé, D., et al. 1997. Cloning of human 2H9 heterogeneous nuclear ribonucleoproteins. Relation with splicing and early heat shock-induced splicing arrest. *J. Biol. Chem.* 272: 1827-1836.
2. Honoré, B. 2000. The hnRNP 2H9 gene, which is involved in the splicing reaction, is a multiply spliced gene. *Biochim. Biophys. Acta* 1492: 108-119.
3. Mahé, D., et al. 2000. Spatiotemporal regulation of hnRNP M and 2H9 gene expression during mouse embryonic development. *Biochim. Biophys. Acta* 1492: 414-424.
4. Yagüe, J., et al. 2000. A post-translational modification of nuclear proteins, N(G),N(G)-dimethyl-Arg, found in a natural HLA class I peptide ligand. *Protein Sci.* 9: 2210-2217.
5. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 602324. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Andersen, J.S., et al. 2005. Nucleolar proteome dynamics. *Nature* 433: 77-83.
7. Olsen, J.V., et al. 2006. Global, *in vivo*, and site-specific phosphorylation dynamics in signaling networks. *Cell* 127: 635-648.

CHROMOSOMAL LOCATION

Genetic locus: HNRNPH3 (human) mapping to 10q21.3.

PRODUCT

hnRNP H3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see hnRNP H3 shRNA Plasmid (h): sc-90762-SH and hnRNP H3 shRNA (h) Lentiviral Particles: sc-90762-V as alternate gene silencing products.

For independent verification of hnRNP H3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-90762A, sc-90762B and sc-90762C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

hnRNP H3 siRNA (h) is recommended for the inhibition of hnRNP H3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

hnRNP H3 (D-4): sc-376416 is recommended as a control antibody for monitoring of hnRNP H3 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor hnRNP H3 gene expression knockdown using RT-PCR Primer: hnRNP H3 (h)-PR: sc-90762-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.