

NDUFB11 siRNA (h): sc-90885

BACKGROUND

Complex I (also known as NADH dehydrogenase) of the electron transport chain (respiratory chain) is an enzymatic complex that catalyzes the transfer of electrons from NADH to ubiquinone. Free energy from the reaction is conserved in the transfer of protons into the intermembrane space to create an electrochemical proton gradient, a driving force for ATP synthesis. Complex I is a complicated, multi-protein, L-shaped complex composed of at least 45 different subunits and located in the mitochondrial inner membrane. NDUFB11 (NADH dehydrogenase (ubiquinone) 1 β subcomplex subunit 11), also known as ESSS, Np15, Np17.3 (neuronal protein 17.3) or p17.3, is a hydrophobic transmembrane protein belonging to the complex I NDUFB11 subunit family. Ubiquitously expressed, NDUFB11 localizes to the inner membrane of the mitochondrion and functions as an accessory subunit of complex I. The cAMP-dependent phosphorylation of NDUFB11 is important for the regulation of complex I activity.

REFERENCES

1. Cui, Y., et al. 1999. Cloning and tissue expressional characterization of a full-length cDNA encoding human neuronal protein P17.3. *Biochem. Genet.* 37: 175-185.
2. Online Mendelian Inheritance in Man, OMIM™. 2002. Johns Hopkins University, Baltimore, MD. MIM Number: 300403. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
3. Janssen, R.J., et al. 2006. Mitochondrial complex I: structure, function and pathology. *J. Inherit. Metab. Dis.* 29: 499-515.
4. Petruzzella, V., et al. 2007. The NDUFB11 gene is not a modifier in Leber hereditary optic neuropathy. *Biochem. Biophys. Res. Commun.* 355: 181-187.
5. Fernandez-Moreira, D., et al. 2007. X-linked NDUF41 gene mutations associated with mitochondrial encephalomyopathy. *Ann. Neurol.* 61: 73-83.
6. Gurok, U., et al. 2007. Expression of NDUFB11 encoding the neuronal protein 15.6 during neurite outgrowth and development. *Gene Expr. Patterns* 7: 370-374.
7. De Rasmio, D., et al. 2008. cAMP-dependent protein kinase regulates the mitochondrial import of the nuclear encoded NDUF41 subunit of complex I. *Cell. Signal.* 20: 989-997.

CHROMOSOMAL LOCATION

Genetic locus: NDUFB11 (human) mapping to Xp11.23.

PRODUCT

NDUFB11 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NDUFB11 shRNA Plasmid (h): sc-90885-SH and NDUFB11 shRNA (h) Lentiviral Particles: sc-90885-V as alternate gene silencing products.

For independent verification of NDUFB11 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-90885A, sc-90885B and sc-90885C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

NDUFB11 siRNA (h) is recommended for the inhibition of NDUFB11 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

NDUFB11 (E-6): sc-393110 is recommended as a control antibody for monitoring of NDUFB11 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NDUFB11 gene expression knockdown using RT-PCR Primer: NDUFB11 (h)-PR: sc-90885-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.