

LHFPL1 siRNA (h): sc-90998

BACKGROUND

LHFPL1 (lipoma HMGIC fusion partner-like 1 protein) is a 220 amino acid protein that widely expressed, with high levels found in the lungs, thymus, skeletal muscles, colon and ovaries. Belonging to the LHP family, LHFPL1 exists as three alternatively spliced isoforms. The gene that encodes LHFPL1 is located on the X chromosome. The X and Y chromosomes are the human sex chromosomes. Chromosome X consists of about 153 million base pairs and nearly 1,000 genes. The combination of an X and Y chromosome lead to normal male development while two copies of X lead to normal female development. There are a number of conditions related to an unusual number and combination of sex chromosomes being inherited, including Turner's syndrome, Klinefelter's syndrome and triple X syndrome. Color blindness, hemophilia, and Duchenne muscular dystrophy are well known X chromosome-linked conditions which affect males more frequently as males carry a single X chromosome.

REFERENCES

1. Gianfrancesco, F., Sanges, R., Esposito, T., Tempesta, S., Rao, E., Rappold, G., Archidiacono, N., Graves, J.A., Forabosco, A. and D'Urso, M. 2001. Differential divergence of three human pseudoautosomal genes and their mouse homologs: implications for sex chromosome evolution. *Genome Res.* 11: 2095-2100.
2. Bernardino-Sgherri, J., Flagiello, D. and Dutrillaux, B. 2002. Overall DNA methylation and chromatin structure of normal and abnormal X chromosomes. *Cytogenet. Genome Res.* 99: 85-91.
3. Huang, C., Guo, J., Liu, S., Shan, Y., Wu, S., Cai, Y. and Yu, L. 2004. Isolation, tissue distribution and prokaryotic expression of a novel human X-linked gene LHFPL1. *DNA Seq.* 15: 299-302.
4. Deeb, S.S. 2005. The molecular basis of variation in human color vision. *Clin. Genet.* 67: 369-377.
5. Bojesen, A., Kristensen, K., Birkebaek, N.H., Fedder, J., Mosekilde, L., Bennett, P., Laurberg, P., Frystyk, J., Flyvbjerg, A., Christiansen, J.S. and Gravholt, C.H. 2006. The metabolic syndrome is frequent in Klinefelter's syndrome and is associated with abdominal obesity and hypogonadism. *Diabetes Care* 29: 1591-1598.
6. Maggio, M.C., Liotta, A., De Grazia, E., Cimador, M., Di Pace, R. and Corsello, G. 2007. Polycystic ovary and gonadoblastoma in Turner's syndrome. *Minerva Pediatr.* 59: 397-401.
7. Helderma-van den Enden, A.T., de Jong, R., den Dunnen, J.T., Houwing-Duistermaat, J.J., Kneppers, A.L., Ginjaar, H.B., Breuning, M.H. and Bakker, E. 2009. Recurrence risk due to germ line mosaicism: Duchenne and Becker muscular dystrophy. *Clin. Genet.* 75: 465-472.

CHROMOSOMAL LOCATION

Genetic locus: LHFPL1 (human) mapping to Xq23.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

LHFPL1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see LHFPL1 shRNA Plasmid (h): sc-90998-SH and LHFPL1 shRNA (h) Lentiviral Particles: sc-90998-V as alternate gene silencing products.

For independent verification of LHFPL1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-90998A, sc-90998B and sc-90998C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

LHFPL1 siRNA (h) is recommended for the inhibition of LHFPL1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor LHFPL1 gene expression knockdown using RT-PCR Primer: LHFPL1 (h)-PR: sc-90998-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.