



## NXT-2 siRNA (h): sc-91277

### BACKGROUND

Protein transport across the nucleus is a selective, multistep process involving several cytoplasmic factors including Ran. Nuclear transport factor 2 (NTF2) regulates Ran function in a non-catalytic fashion and mediates Ran-GDP targeting to the nucleus. Nucleotide-dependent conformations of Ran alter the site of interaction that would otherwise permit the binding of NTF2 to Ran-GTP. A member of the NXT family, NXT-2 (NTF2-related export protein 2) is a 142 amino acid protein containing an NTF2 domain. NXT-2 shuttles between the nucleus and the cytoplasm, regulating protein export for NES-containing proteins. NXT-2 also plays a role in mRNA nuclear export and associates with TAP, NXF2, NXF3 and NXF5. Four isoforms are produced by alternative splicing events.

### REFERENCES

1. Smith, A., Brownawell, A. and Macara, I.G. 1998. Nuclear import of Ran is mediated by the transport factor NTF2. *Curr. Biol.* 8: 1403-1406.
2. Ribbeck, K., Lipowsky, G., Kent, H.M., Stewart, M. and Görlich, D. 1998. NTF2 mediates nuclear import of Ran. *EMBO J.* 17: 6587-6598.
3. Black, B.E., Levesque, L., Holaska, J.M., Wood, T.C. and Paschal, B.M. 1999. Identification of an NTF2-related factor that binds Ran-GTP and regulates nuclear protein export. *Mol. Cell. Biol.* 19: 8616-8624.
4. Ossareh-Nazari, B., Maison, C., Black, B.E., Levesque, L., Paschal, B.M. and Dargemont, C. 2000. RanGTP-binding protein NXT1 facilitates nuclear export of different classes of RNA *in vitro*. *Mol. Cell. Biol.* 20: 4562-4571.
5. Herold, A., Suyama, M., Rodrigues, J.P., Braun, I.C., Kutay, U., Carmo-Fonseca, M., Bork, P. and Izaurralde, E. 2000. TAP (NXF1) belongs to a multigene family of putative RNA export factors with a conserved modular architecture. *Mol. Cell. Biol.* 20: 8996-9008.
6. Online Mendelian Inheritance in Man, OMIM™. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 300320. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/300320>
7. Huang, H., Zhang, B., Hartenstein, P.A., Chen, J.N. and Lin, S. 2005. NXT2 is required for embryonic heart development in zebrafish. *BMC Dev. Biol.* 5: 7.

### CHROMOSOMAL LOCATION

Genetic locus: NXT2 (human) mapping to Xq23.

### PRODUCT

NXT-2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see NXT-2 shRNA Plasmid (h): sc-91277-SH and NXT-2 shRNA (h) Lentiviral Particles: sc-91277-V as alternate gene silencing products.

For independent verification of NXT-2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-91277A, sc-91277B and sc-91277C.

### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

### APPLICATIONS

NXT-2 siRNA (h) is recommended for the inhibition of NXT-2 expression in human cells.

### SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

### RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor NXT-2 gene expression knockdown using RT-PCR Primer: NXT-2 (h)-PR: sc-91277-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

### RESEARCH USE

For research use only, not for use in diagnostic procedures.

### PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.