

# MAGE-B10 siRNA (h): sc-91279

## BACKGROUND

The melanoma-associated antigen (MAGE) family consists of a number of antigens recognized by cytotoxic T lymphocytes. The MAGE genes were initially isolated from different kinds of tumors and, based on their virtually exclusive tumor-specific expression in adult tissues, they have been used as targets for cancer immunotherapy. MAGE genes encode for tumor-rejection antigens that are expressed in tumors of different histologic types and in normal testis and placenta. MAGE-B10 (melanoma-associated antigen B10) is a 347 amino acid protein that contains one MAGE domain. The gene encoding MAGE-B10 maps to human chromosome X, which consists of about 153 million base pairs and nearly 1,000 genes. Color blindness, hemophilia and Duchenne muscular dystrophy are well known X chromosome-linked conditions which affect males more frequently, as males carry a single X chromosome.

## REFERENCES

1. Chomez, P., De Backer, O., Bertrand, M., De Plaen, E., Boon, T. and Lucas, S. 2001. An overview of the MAGE gene family with the identification of all human members of the family. *Cancer Res.* 61: 5544-5551.
2. Kirkin, A.F., Dzhandzhugazyan, K.N. and Zeuthen, J. 2002. Cancer/testis antigens: structural and immunobiological properties. *Cancer Invest.* 20: 222-236.
3. Deeb, S.S. 2005. The molecular basis of variation in human color vision. *Clin. Genet.* 67: 369-377.
4. Helderma-van den Enden, A.T., de Jong, R., den Dunnen, J.T., Houwing-Duistermaat, J.J., Kneppers, A.L., Ginjaar, H.B., Breuning, M.H. and Bakker, E. 2009. Recurrence risk due to germ line mosaicism: Duchenne and Becker muscular dystrophy. *Clin. Genet.* 75: 465-472.
5. Online Mendelian Inheritance in Man, OMIM<sup>™</sup>. 2009. Johns Hopkins University, Baltimore, MD. MIM Number: 300761. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Kasper, C.K. and Buzin, C.H. 2009. Mosaicism and haemophilia. *Haemophilia*. E-published.
7. Marcar, L., MacLaine, N.J., Hupp, T.R. and Meek, D.W. 2010. Mage-A cancer/testis antigens inhibit p53 function by blocking its interaction with chromatin. *Cancer Res.* 70: 10362-10370.
8. Sang, M., Wang, L., Ding, C., Zhou, X., Wang, B., Wang, L., Lian, Y. and Shan, B. 2011. Melanoma-associated antigen genes-an update. *Cancer Lett.* 302: 85-90.
9. Feng, Y., Gao, J. and Yang, M. 2011. When MAGE meets RING: insights into biological functions of MAGE proteins. *Protein Cell* 2: 7-12.

## CHROMOSOMAL LOCATION

Genetic locus: MAGEB10 (human) mapping to Xp21.3.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## PRODUCT

MAGE-B10 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see MAGE-B10 shRNA Plasmid (h): sc-91279-SH and MAGE-B10 shRNA (h) Lentiviral Particles: sc-91279-V as alternate gene silencing products.

For independent verification of MAGE-B10 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-91279A, sc-91279B and sc-91279C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

MAGE-B10 siRNA (h) is recommended for the inhibition of MAGE-B10 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor MAGE-B10 gene expression knockdown using RT-PCR Primer: MAGE-B10 (h)-PR: sc-91279-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.