

RHOBTB3 siRNA (h): sc-91688

BACKGROUND

The Rho subfamily of Ras-related GTPases controls multiple aspects of cell function, including cytoskeletal rearrangement, nuclear signaling and cell growth. RHOBTB1 (Rho-related BTB domain-containing protein 1) and RHOBTB3 (Rho-related BTB domain-containing protein 3) each contain two BTB (POZ) domains and belong to the RhoBTB subfamily of Rho GTPases. Members of the RhoBTB subfamily are evolutionarily conserved and are characterized by a proline-rich region, a GTPase domain and two tandem BTB repeats. While both RHOBTB1 and RHOBTB3 are expressed ubiquitously, RHOBTB1 is found at high levels in placenta, stomach, testis, kidney and skeletal muscle, whereas RHOBTB3 is found at high levels in neural and cardiac tissues. RHOBTB1 is thought to play a role in GTPase-mediated signaling and may participate in organization of the Actin filament system. Additionally, RHOBTB1 expression is decreased in head and neck carcinomas, suggesting a possible role for RHOBTB1 as a tumor suppressor.

REFERENCES

1. Rivero, F., Dislich, H., Glöckner, G. and Noegel, A.A. 2001. The *Dictyostelium discoideum* family of Rho-related proteins. *Nucleic Acids Res.* 29: 1068-1079.
2. Ramos, S., Khademi, F., Somesh, B.P. and Rivero, F. 2002. Genomic organization and expression profile of the small GTPases of the RhoBTB family in human and mouse. *Gene* 298: 147-157.
3. Aspenström, P., Fransson, A. and Saras, J. 2004. Rho GTPases have diverse effects on the organization of the Actin filament system. *Biochem. J.* 377: 327-337.
4. Vlahou, G. and Rivero, F. 2006. Rho GTPase signaling in *Dictyostelium discoideum*: insights from the genome. *Eur. J. Cell Biol.* 85: 947-959.
5. Beder, L.B., Gunduz, M., Ouchida, M., Gunduz, E., Sakai, A., Fukushima, K., Nagatsuka, H., Ito, S., Honjo, N., Nishizaki, K. and Shimizu, K. 2006. Identification of a candidate tumor suppressor gene RHOBTB1 located at a novel allelic loss region 10q21 in head and neck cancer. *J. Cancer Res. Clin. Oncol.* 132: 19-27.
6. Aspenström, P., Ruusala, A. and Pacholsky, D. 2007. Taking Rho GTPases to the next level: the cellular functions of atypical Rho GTPases. *Exp. Cell Res.* 313: 3673-3679.
7. Boureux, A., Vignal, E., Faure, S. and Fort, P. 2007. Evolution of the Rho family of Ras-like GTPases in eukaryotes. *Mol. Biol. Evol.* 24: 203-216.
8. Berthold, J., Schenkova, K. and Rivero, F. 2008. Rho GTPases of the RhoBTB subfamily and tumorigenesis. *Acta Pharmacol. Sin.* 29: 285-295.

CHROMOSOMAL LOCATION

Genetic locus: RHOBTB3 (human) mapping to 5q15.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

RHOBTB3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RHOBTB3 shRNA Plasmid (h): sc-91688-SH and RHOBTB3 shRNA (h) Lentiviral Particles: sc-91688-V as alternate gene silencing products.

For independent verification of RHOBTB3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-91688A, sc-91688B and sc-91688C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RHOBTB3 siRNA (h) is recommended for the inhibition of RHOBTB3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RESEARCH USE

For research use only, not for use in diagnostic procedures.