



DMXL1 siRNA (h): sc-91728

BACKGROUND

WD-repeats are motifs that are found in a variety of proteins and are characterized by a conserved core of 40-60 amino acids that commonly form a tertiary propeller structure. While proteins that contain WD-repeats participate in a wide range of cellular functions, they are generally involved in regulatory mechanisms concerning chromatin assembly, cell cycle control, signal transduction, RNA processing, apoptosis and vesicular trafficking. DMXL1 (DmX-like protein 1), also known as XL1 or FLJ44772, is a 3,027 amino acid protein that contains 15 WD repeats. DMXL1 is expressed in foreskin, tonsils, parathyroid, breast, eye, heart, bone, testis, small intestine, uterus and placenta. The gene that encodes DMXL1 maps to human chromosome 5.

REFERENCES

1. Komachi, K., Redd, M.J. and Johnson, A.D. 1994. The WD repeats of Tup1 interact with the homeo domain protein α 2. *Genes Dev.* 8: 2857-2867.
2. Kolman, M.F. and Egelhoff, T.T. 1997. *Dictyostelium* myosin heavy chain kinase A subdomains. Coiled-coil and wd repeat roles in oligomerization and substrate targeting. *J. Biol. Chem.* 272: 16904-16910.
3. Wolf, D.A., McKeon, F. and Jackson, P.K. 1999. F-box/WD-repeat proteins pop1p and Sud1p/Pop2p form complexes that bind and direct the proteolysis of cdc18p. *Curr. Biol.* 9: 373-376.
4. Suprenant, K.A., Tuxhorn, J.A., Daggett, M.A., Ahrens, D.P., Hostetler, A., Palange, J.M., VanWinkle, C.E. and Livingston, B.T. 2000. Conservation of the WD-repeat, microtubule-binding protein, EMAP, in sea urchins, humans, and the nematode *C. elegans*. *Dev. Genes Evol.* 210: 2-10.
5. Kraemer, C., Enklaar, T., Zabel, B. and Schmidt, E.R. 2000. Mapping and structure of DMXL1, a human homologue of the DmX gene from *Drosophila melanogaster* coding for a WD repeat protein. *Genomics* 64: 97-101.
6. Hisbergues, M., Gaitatzes, C.G., Joset, F., Bedu, S. and Smith, T.F. 2001. A noncanonical WD-repeat protein from the cyanobacterium *Synechocystis* PCC6803: structural and functional study. *Protein Sci.* 10: 293-300.

CHROMOSOMAL LOCATION

Genetic locus: DMXL1 (human) mapping to 5q23.1.

PRODUCT

DMXL1 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see DMXL1 shRNA Plasmid (h): sc-91728-SH and DMXL1 shRNA (h) Lentiviral Particles: sc-91728-V as alternate gene silencing products.

For independent verification of DMXL1 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-91728A, sc-91728B and sc-91728C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

DMXL1 siRNA (h) is recommended for the inhibition of DMXL1 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor DMXL1 gene expression knockdown using RT-PCR Primer: DMXL1 (h)-PR: sc-91728-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.