

BTNL3 siRNA (h): sc-91733

BACKGROUND

Butyrophilin is a glycoprotein that is specifically expressed on the apical surface of mammary epithelial cells during lactation and becomes incorporated as an integral protein into the membrane of the milk fat globule during the budding and secretion of fat droplets into milk. BTNL3 (butyrophilin-like protein 3), also known as BTNLR (butyrophilin-like receptor) or COLF4100, is a 466 amino acid single-pass type I membrane protein that localizes to the outer membrane of milk fat-secreting droplets of mammary epithelial cells. BTNL3 is a member of the immunoglobulin superfamily and BTN/MOG family, and exists as two alternatively spliced isoforms that are encoded by a gene located on human chromosome 5. BTNL3 contains one B30.2/SPRY domain and a single Ig-like V-type (immunoglobulin-like) domain.

REFERENCES

1. Mather, I.H., et al. 1993. A review of the molecular and cellular biology of butyrophilin, the major protein of bovine milk fat globule membrane. *J. Dairy Sci.* 76: 3832-3850.
2. Ogg, S.L., et al. 1996. Structural organization and mammary-specific expression of the butyrophilin gene. *Mamm. Genome* 7: 900-905.
3. Shibui, A., et al. 1999. Cloning, expression analysis, and chromosomal localization of a novel butyrophilin-like receptor. *J. Hum. Genet.* 44: 249-252.
4. Stammers, M., et al. 2000. BTL-II: a polymorphic locus with homology to the butyrophilin gene family, located at the border of the major histocompatibility complex class II and class III regions in human and mouse. *Immunogenetics* 51: 373-382.
5. Online Mendelian Inheritance in Man, OMIM™. 2001. Johns Hopkins University, Baltimore, MD. MIM Number: 606192. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Cavaletto, M., et al. 2002. A proteomic approach to evaluate the butyrophilin gene family expression in human milk fat globule membrane. *Proteomics* 2: 850-856.

CHROMOSOMAL LOCATION

Genetic locus: BTNL3 (human) mapping to 5q35.3.

PRODUCT

BTNL3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see BTNL3 shRNA Plasmid (h): sc-91733-SH and BTNL3 shRNA (h) Lentiviral Particles: sc-91733-V as alternate gene silencing products.

For independent verification of BTNL3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-91733A, sc-91733B and sc-91733C.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

BTNL3 siRNA (h) is recommended for the inhibition of BTNL3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor BTNL3 gene expression knockdown using RT-PCR Primer: BTNL3 (h)-PR: sc-91733-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.