

OATP-I siRNA (h): sc-91933

BACKGROUND

The organic anion transporter family of proteins mediate hepatic uptake of cardiac glycosides. OATP-I (organic anion transporter F), also known as SLC06A1 (solute carrier organic anion transporter family member 6A1), gonad-specific transporter (GST), cancer/testis antigen 48 (CT48) or SLC21A14 (solute carrier family 21 member 19), is a 719 amino acid member of the organic anion transporter protein family. As a multi-pass membrane protein, OATP-I is thought to mediate the Na⁺-independent transport of organic anions. OATP-I is highly expressed in testis, with lower expression found in brain, placenta and spleen. OATP-I is expressed as three isoforms produced by alternative splicing events.

REFERENCES

1. Hsiang, B., Zhu, Y., Wang, Z., Wu, Y., Sasseville, V., Yang, W.P. and Kirchgesner, T.G. 1999. A novel human hepatic organic anion transporting polypeptide (OATP2). *J. Biol. Chem.* 274: 37161-37168.
2. König, J., Cui, Y., Nies, A.T. and Keppler, D. 2000. Localization and genomic organization of a new hepatocellular organic anion transporting polypeptide. *J. Biol. Chem.* 275: 23161-23168.
3. Cai, S.Y., Wang, W., Soroka, C.J., Ballatori, N. and Boyer, J.L. 2002. An evolutionarily ancient Oatp: insights into conserved functional domains of these proteins. *Am. J. Physiol. Gastrointest. Liver Physiol.* 282: G702-G710.
4. Pizzagalli, F., Hagenbuch, B., Stieger, B., Klenk, U., Folkers, G. and Meier, P.J. 2002. Identification of a novel human organic anion transporting polypeptide as a high affinity thyroxine transporter. *Mol. Endocrinol.* 16: 2283-2296.
5. Patel, P., Weerasekera, N., Hitchins, M., Boyd, C.A., Johnston, D.G. and Williamson, C. 2002. Semi quantitative expression analysis of Mdr-3, FIC1, BSEP, OATP-A, OATP-C, OATP-D, OATP-E and NTCP gene transcripts in 1st and 3rd trimester human placenta. *Placenta* 24: 39-44.
6. Suzuki, T., Onogawa, T., Asano, N., Mizutamari, H., Mikkaichi, T., Tanemoto, M., Abe, M., Satoh, F., Unno, M., Nunoki, K., Suzuki, M., Hishinuma, T., Goto, J., Shimosegawa, T., Matsuno, S., et al. 2003. Identification and characterization of novel rat and human gonad-specific organic anion transporters. *Mol. Endocrinol.* 17: 1203-1215.
7. Hagenbuch, B. and Meier, P.J. 2004. Organic anion transporting polypeptides of the OATP/SLC21 family: phylogenetic classification as OATP/SLC20 superfamily, new nomenclature and molecular/functional properties. *Pflügers Arch.* 447: 653-665.
8. König, J., Seithel, A., Gradhand, U. and Fromm, M.F. 2006. Pharmacogenomics of human OATP transporters. *Naunyn Schmiedeberg's Arch. Pharmacol.* 372: 432-443.
9. Seithel, A., Glaeser, H., Fromm, M.F. and König, J. 2008. The functional consequences of genetic variations in transporter genes encoding human organic anion-transporting polypeptide family members. *Expert Opin. Drug Metab. Toxicol.* 4: 51-64.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

CHROMOSOMAL LOCATION

Genetic locus: SLC06A1 (human) mapping to 5q21.1.

PRODUCT

OATP-I siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see OATP-I shRNA Plasmid (h): sc-91933-SH and OATP-I shRNA (h) Lentiviral Particles: sc-91933-V as alternate gene silencing products.

For independent verification of OATP-I (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-91933A, sc-91933B and sc-91933C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

OATP-I siRNA (h) is recommended for the inhibition of OATP-I expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor OATP-I gene expression knockdown using RT-PCR Primer: OATP-I (h)-PR: sc-91933-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.