

ARRDC3 siRNA (h): sc-92033

BACKGROUND

ARRDC3 (arrestin domain containing 3), also known as TLIMP, is a 414 amino acid protein that belongs to the arrestin family and is encoded by a gene which maps to human chromosome 5q14.3. With 181 million base pairs encoding around 1,000 genes, chromosome 5 comprises over 6% of human genomic DNA. Chromosome 5 is associated with Cockayne syndrome through the ERCC8 gene, as well as with familial adenomatous polyposis through the adenomatous polyposis coli (APC) tumor suppressor gene. Treacher Collins syndrome is also chromosome 5-associated and is caused by insertions or deletions within the TCOF1 gene. Deletion of the p arm of chromosome 5 leads to Cri du chat syndrome, while deletion of 5q or chromosome 5 altogether is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

REFERENCES

1. Dixon, M.J., et al. 1991. The gene for Treacher Collins syndrome maps to the long arm of chromosome 5. *Am. J. Hum. Genet.* 49: 17-22.
2. Joslyn, G., et al. 1991. Identification of deletion mutations and three new genes at the familial polyposis locus. *Cell* 66: 601-613.
3. Kinzler, K.W., et al. 1991. Identification of FAP locus genes from chromosome 5q21. *Science* 253: 661-665.
4. Nishisho, I., et al. 1991. Mutations of chromosome 5q21 genes in FAP and colorectal cancer patients. *Science* 253: 665-669.
5. Prieschl, E.E., et al. 1996. The murine homolog of TB2/DP1, a gene of the familial adenomatous polyposis (FAP) locus. *Gene* 169: 215-218.
6. Nagase, T., et al. 2000. Prediction of the coding sequences of unidentified human genes. XVI. The complete sequences of 150 new cDNA clones from brain which code for large proteins *in vitro*. *DNA Res.* 7: 65-73.
7. Oka, S., et al. 2006. Thioredoxin-binding protein-2-like inducible membrane protein is a novel vitamin D₃ and peroxisome proliferator-activated receptor (PPAR) γ ligand target protein that regulates PPAR γ signaling. *Endocrinology* 147: 733-743.
8. Shin, S.M., et al. 2006. HCCR-1-interacting molecule "deleted in polyposis 1" plays a tumor-suppressor role in colon carcinogenesis. *Gastroenterology* 130: 2074-2086.

CHROMOSOMAL LOCATION

Genetic locus: ARRDC3 (human) mapping to 5q14.3.

PRODUCT

ARRDC3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ARRDC3 shRNA Plasmid (h): sc-92033-SH and ARRDC3 shRNA (h) Lentiviral Particles: sc-92033-V as alternate gene silencing products.

For independent verification of ARRDC3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-92033A, sc-92033B and sc-92033C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ARRDC3 siRNA (h) is recommended for the inhibition of ARRDC3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ARRDC3 gene expression knockdown using RT-PCR Primer: ARRDC3 (h)-PR: sc-92033-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.