



# SMA5 siRNA (h): sc-92040

## BACKGROUND

SMA5 is a 140 amino acid protein that is encoded by a gene mapping to human chromosome 5, which contains 181 million base pairs and comprises nearly 6% of the human genome. Chromosome 5 is associated with Cockayne syndrome through the ERCC8 gene and familial adenomatous polyposis through the adenomatous polyposis coli (APC) tumor suppressor gene. Treacher Collins syndrome is also chromosome 5-associated and is caused by insertions or deletions within the TCOF1 gene. Deletion of the p arm of chromosome 5 leads to Cri du chat syndrome, while deletion of the q arm of chromosome 5 altogether is common in therapy-related acute myelogenous leukemias and myelodysplastic syndrome.

## REFERENCES

1. Rauch, A. and Dörr, H.G. 2007. Chromosome 5q subtelomeric deletion syndrome. *Am. J. Med. Genet. C Semin. Med. Genet.* 145C: 372-376.
2. Villa, N., Redaelli, S., Borroni, C., Colombo, C., Roncaglia, N., Sala, E., Crosti, F., Cappellini, A. and Dalprà, L. 2007. Fetal trisomy 5 mosaicism: case report and literature review. *Am. J. Med. Genet. A* 143A: 2343-2346.
3. Shadduck, R.K., Latsko, J.M., Rossetti, J.M., Haq, B. and Abdulhaq, H. 2007. Recent advances in myelodysplastic syndromes. *Exp. Hematol.* 35: 137-143.
4. Falini, B., Nicoletti, I., Bolli, N., Martelli, M.P., Liso, A., Gorello, P., Mandelli, F., Mecucci, C. and Martelli, M.F. 2007. Translocations and mutations involving the nucleophosmin (NPM1) gene in lymphomas and leukemias. *Haematologica* 92: 519-532.
5. Kristoffersen, K.E. 2008. Speech and language development in cri du chat syndrome: a critical review. *Clin. Linguist. Phon.* 22: 443-457.
6. Valent, P. 2008. Revealing the pathogenesis of the 5q- syndrome. *Eur. J. Clin. Invest.* 38: 539-540.
7. Buysse, K., Crepel, A., Menten, B., Pattyn, F., Antonacci, F., Veltman, J.A., Larsen, L.A., Tümer, Z., de Klein, A., van de Laar, I., Devriendt, K., Mortier, G. and Speleman, F. 2008. Mapping of 5q35 chromosomal rearrangements within a genomically unstable region. *J. Med. Genet.* 45: 672-678.
8. Azman, B.Z., Akhir, S.M., Zilfalil, B.A. and Ankathil, R. 2008. Two cases of deletion 5p syndrome: one with paternal involvement and another with atypical presentation. *Singapore Med. J.* 49: e98-e100.

## CHROMOSOMAL LOCATION

Genetic locus: SMA5 (human) mapping to 5p15.33.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## PRODUCT

SMA5 siRNA (h) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SMA5 shRNA Plasmid (h): sc-92040-SH and SMA5 shRNA (h) Lentiviral Particles: sc-92040-V as alternate gene silencing products.

For independent verification of SMA5 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-92040A and sc-92040B.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

SMA5 siRNA (h) is recommended for the inhibition of SMA5 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SMA5 gene expression knockdown using RT-PCR Primer: SMA5 (h)-PR: sc-92040-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.