



Ribosomal Protein S29 siRNA (h): sc-92092

BACKGROUND

Ribosomes, the organelles that catalyze protein synthesis, are composed of a small subunit (40S) and a large subunit (60S) that consist of over 80 distinct ribosomal proteins. Mammalian ribosomal proteins are encoded by multigene families that contain processed pseudogenes and one functional intron-containing gene within their coding regions. Ribosomal Protein S29, also known as RPS29, is a 56 amino acid ribosomal protein that exists as a component of the 40S subunit and contains one C2-C2 zinc finger-like domain. Localized to the cytoplasm, Ribosomal Protein S29 binds zinc as a cofactor and is thought to enhance the tumor suppressor activity of Rap 1A, possibly playing an indirect role in tumor suppression. Like most ribosomal proteins, Ribosomal Protein S29 exists as multiple processed pseudogenes that are scattered throughout the genome.

REFERENCES

1. Wool, I.G., Chan, Y.L. and Glück, A. 1995. Structure and evolution of mammalian ribosomal proteins. *Biochem. Cell Biol.* 73: 933-947.
2. Frigerio, J.M., Dagorn, J.C. and Iovanna, J.L. 1995. Cloning, sequencing and expression of the L5, L21, L27a, L28, S5, S9, S10 and S29 human ribosomal protein mRNAs. *Biochim. Biophys. Acta* 1262: 64-68.
3. Kondoh, N., Noda, M., Fisher, R.J., Schweinfest, C.W., Papas, T.S., Kondoh, A., Samuel, K.P. and Oikawa, T. 1996. The Ribosomal Protein S29 increases tumor suppressor activity of K rev-1 gene on v-K ras-transformed NIH/3T3 cells. *Biochim. Biophys. Acta* 1313: 41-46.
4. Vladimirov, S.N., Ivanov, A.V., Karpova, G.G., Musolyamov, A.K., Egorov, T.A., Thiede, B., Wittmann-Liebold, B. and Otto, A. 1996. Characterization of the human small-ribosomal-subunit proteins by N-terminal and internal sequencing, and mass spectrometry. *Eur. J. Biochem.* 239: 144-149.
5. Kenmochi, N., Kawaguchi, T., Rozen, S., Davis, E., Goodman, N., Hudson, T.J., Tanaka, T. and Page, D.C. 1998. A map of 75 human ribosomal protein genes. *Genome Res.* 8: 509-523.
6. Online Mendelian Inheritance in Man, OMIM™. 1999. Johns Hopkins University, Baltimore, MD. MIM Number: 603633. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
7. Coppock, D., Kopman, C., Gudas, J. and Cina-Poppe, D.A. 2000. Regulation of the quiescence-induced genes: quiescins Q6, Decorin, and Ribosomal Protein S29. *Biochem. Biophys. Res. Commun.* 269: 604-610.

CHROMOSOMAL LOCATION

Genetic locus: RPS29 (human) mapping to 14q21.3.

PRODUCT

Ribosomal Protein S29 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Ribosomal Protein S29 shRNA Plasmid (h): sc-92092-SH and Ribosomal Protein S29 shRNA (h) Lentiviral Particles: sc-92092-V as alternate gene silencing products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Ribosomal Protein S29 siRNA (h) is recommended for the inhibition of Ribosomal Protein S29 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

GENE EXPRESSION MONITORING

Ribosomal Protein S29 (3G9): sc-517071 is recommended as a control antibody for monitoring of Ribosomal Protein S29 gene expression knockdown by Western Blotting (starting dilution 1:200, dilution range 1:100-1:1000) or immunofluorescence (starting dilution 1:50, dilution range 1:50-1:500).

To ensure optimal results, the following support reagents are recommended: 1) Western Blotting: use m-IgG κ BP-HRP: sc-516102 or m-IgG κ BP-HRP (Cruz Marker): sc-516102-CM (dilution range: 1:1000-1:10000), Cruz Marker™ Molecular Weight Standards: sc-2035, UltraCruz® Blocking Reagent: sc-516214 and Western Blotting Luminol Reagent: sc-2048. 2) Immunofluorescence: use m-IgG κ BP-FITC: sc-516140 or m-IgG κ BP-PE: sc-516141 (dilution range: 1:50-1:200) with UltraCruz® Mounting Medium: sc-24941 or UltraCruz® Hard-set Mounting Medium: sc-359850.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Ribosomal Protein S29 gene expression knockdown using RT-PCR Primer: Ribosomal Protein S29 (h)-PR: sc-92092-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.