GPR137C siRNA (h): sc-92110



The Power to Question

BACKGROUND

G protein-coupled receptors (GPRs or GPCRs), also known as seven transmembrane receptors, heptahelical receptors or 7TM receptors, are members of the largest protein family and play a role in many different stimulus-response pathways. G protein-coupled receptors mediate extracellular signals into intracellular signals (G protein activation). They respond to a wide variety of signaling molecules, including hormones, neurotransmitters and other proteins and peptides. GPR proteins are usually integral seven pass membrane proteins with some conserved amino acid regions. GPR137C (G protein-coupled receptor 137C), also known as TM7SF1L2 (transmembrane 7 superfamily member 1-like 2 protein), is a 429 amino acid multi-pass membrane protein that belongs to the GPR137 family. The gene encoding GPR137C maps to human chromosome 14q22.1.

REFERENCES

- Fredriksson, R., et al. 2002. Novel human G protein-coupled receptors with long N-terminals containing GPS domains and Ser/Thr-rich regions. FEBS Lett. 531: 407-414.
- 2. Vassilatis, D.K., et al. 2003. The G protein-coupled receptor repertoires of human and mouse. Proc. Natl. Acad. Sci. USA 100: 4903-4908.
- 3. Bjarnadóttir, T.K., et al. 2004. The human and mouse repertoire of the adhesion family of G-protein-coupled receptors. Genomics 84: 23-33.
- 4. Ross, M.T., et al. 2005. The DNA sequence of the human X chromosome. Nature 434: 325-337.
- 5. Bjarnadóttir, T.K., et al. 2007. Identification of novel splice variants of Adhesion G protein-coupled receptors. Gene 387: 38-48.
- Leja, J., et al. 2009. Novel markers for enterochromaffin cells and gastrointestinal neuroendocrine carcinomas. Mod. Pathol. 22: 261-272.

CHROMOSOMAL LOCATION

Genetic locus: GPR137C (human) mapping to 14q22.1.

PRODUCT

GPR137C siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GPR137C shRNA Plasmid (h): sc-92110-SH and GPR137C shRNA (h) Lentiviral Particles: sc-92110-V as alternate gene silencing products.

For independent verification of GPR137C (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-92110A, sc-92110B and sc-92110C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20 $^{\circ}$ C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20 $^{\circ}$ C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNAse-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNAse-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GPR137C siRNA (h) is recommended for the inhibition of GPR137C expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GPR137C gene expression knockdown using RT-PCR Primer: GPR137C (h)-PR: sc-92110-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

Santa Cruz Biotechnology, Inc. 1.800.457.3801 831.457.3801 fax 831.457.3801 Europe +00800 4573 8000 49 6221 4503 0 www.scbt.com