

RNase 2 siRNA (h): sc-92235

BACKGROUND

RNase 2 [ribonuclease, RNase A family, 2 (liver, eosinophil-derived neurotoxin)], also known as non-secretory ribonuclease, EDN (eosinophil-derived neurotoxin), RNase Upl-2 or RNS2, is a 161 amino acid protein that belongs to the pancreatic ribonuclease family. Localizing to lysosome and cytoplasmic granules, RNase 2 is expressed in leukocytes, liver, spleen, lung and body fluids. RNase 2 functions as a pyrimidine specific nuclease, and has a slight preference for uracil. RNase 2 is capable of various biological activities, including mediation of chemotactic activity and endonucleolytic cleavage of nucleoside 3'-phosphates and 3'-phosphooligonucleotides. The gene encoding RNase 2 maps to human chromosome 14q11.2.

REFERENCES

1. Yasuda, T., Sato, W., Mizuta, K. and Kishi, K. 1988. Genetic polymorphism of human serum ribonuclease I (RNase I). *Am. J. Hum. Genet.* 42: 608-614.
2. Rosenberg, H.F., Tenen, D.G. and Ackerman, S.J. 1989. Molecular cloning of the human eosinophil-derived neurotoxin: a member of the ribonuclease gene family. *Proc. Natl. Acad. Sci. USA* 86: 4460-4464.
3. Hamann, K.J., Ten, R.M., Loegering, D.A., Jenkins, R.B., Heise, M.T., Schad, C.R., Pease, L.R., Gleich, G.J. and Barker, R.L. 1990. Structure and chromosome localization of the human eosinophil-derived neurotoxin and eosinophil cationic protein genes: evidence for intronless coding sequences in the ribonuclease gene superfamily. *Genomics* 7: 535-546.
4. Mastrianni, D.M., Eddy, R.L., Rosenberg, H.F., Corrette, S.E., Shows, T.B., Tenen, D.G. and Ackerman, S.J. 1992. Localization of the human eosinophil Charcot-Leyden crystal protein (lysophospholipase) gene (CLC) to chromosome 19 and the human ribonuclease 2 (eosinophil-derived neurotoxin) and ribonuclease 3 (eosinophil cationic protein) genes (RNS2 and RNS3) to chromosome 14. *Genomics* 13: 240-242.
5. Zhang, J., Dyer, K.D. and Rosenberg, H.F. 2002. RNase 8, a novel RNase A superfamily ribonuclease expressed uniquely in placenta. *Nucleic Acids Res.* 30: 1169-1175.
6. Benner, S.A. 2002. The past as the key to the present: resurrection of ancient proteins from eosinophils. *Proc. Natl. Acad. Sci. USA* 99: 4760-4761.
7. Zhang, J. and Rosenberg, H.F. 2002. Complementary advantageous substitutions in the evolution of an antiviral RNase of higher primates. *Proc. Natl. Acad. Sci. USA* 99: 5486-5491.
8. Online Mendelian Inheritance in Man, OMIM[™]. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 131410. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>

CHROMOSOMAL LOCATION

Genetic locus: RNASE2 (human) mapping to 14q11.2.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

RNase 2 siRNA (h) is a pool of 2 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see RNase 2 shRNA Plasmid (h): sc-92235-SH and RNase 2 shRNA (h) Lentiviral Particles: sc-92235-V as alternate gene silencing products.

For independent verification of RNase 2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-92235A and sc-92235B.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

RNase 2 siRNA (h) is recommended for the inhibition of RNase 2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor RNase 2 gene expression knockdown using RT-PCR Primer: RNase 2 (h)-PR: sc-92235-PR (20 μ l, 561 bp). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.