

ZNF169 siRNA (h): sc-92480

BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. As a member of the Krüppel C₂H₂-type zinc-finger protein family, ZNF169 (zinc finger protein 169) is a 603 amino acid nuclear protein that contains one KRAB domain and thirteen C₂H₂-type zinc fingers. ZNF169 is highly expressed in kidney and weakly expressed in spleen, liver, small intestine and heart, where it functions as a transcription regulator. The gene encoding ZNF169 maps to a region of human chromosome 9q22.32, which has been associated with many human diseases such as colon cancer, migraine auras, basal cell carcinoma, Gorlin syndrome and Extraskeletal myxoid chondrosarcoma.

REFERENCES

1. Levanat, S., Chidambaram, A., Wicking, C., Bray-Ward, P., Pressman, C., Toftgard, R., Gailani, M.R., Myers, J.C., Wainwright, B., Dean, M. and Bale, A.E. 1997. Pulsed-field gel electrophoresis and FISH mapping of chromosome 9q22: placement of a novel zinc finger gene within the NBCCS and ESS1 region. *Cytogenet. Cell Genet.* 76: 208-213.
2. Chidambaram, A., Gailani, M., Gerrard, B., Stewart, C., Goldstein, A., Chumakov, I., Bale, A.E. and Dean, M. 1997. Characterization of a YAC contig containing the NBCCS locus and a novel Krüppel-type zinc finger sequence on chromosome segment 9q22.3. *Genes Chromosomes Cancer* 18: 212-218.
3. Hisaoka, M. and Hashimoto, H. 2005. Extraskeletal myxoid chondrosarcoma: updated clinicopathological and molecular genetic characteristics. *Pathol. Int.* 55: 453-463.
4. Bose, S., Morgan, L.J., Booth, D.R., Goudie, D.R., Ferguson-Smith, M.A. and Richards, F.M. 2006. The elusive multiple self-healing squamous epithelioma (MSSE) gene: further mapping, analysis of candidates, and loss of heterozygosity. *Oncogene* 25: 806-812.
5. Online Mendelian Inheritance in Man, OMIM[™]. 2008. Johns Hopkins University, Baltimore, MD. MIM Number: 603404. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Mosterd, K., Sommer, A., van Marion, A., Lacko, M., Herbergs, J., de Bondt, B.J., van Steensel, M.A. and Kelleners-Smeets, N.W. 2009. Destructive basal cell carcinoma in a patient with basal cell nevus syndrome and an interstitial deletion of chromosome 9q22. *Dermatol. Surg.* 35: 2051-2053.
7. Musani, V., Cretnik, M., Situm, M., Basta-Juzbasic, A. and Levanat, S. 2009. Gorlin syndrome patient with large deletion in 9q22.32-q22.33 detected by quantitative multiplex fluorescent PCR. *Dermatology* 219: 111-118.
8. Gray-McGuire, C., Guda, K., Adrianto, I., Lin, C.P., Natale, L., Potter, J.D., Newcomb, P., Poole, E.M., Ulrich, C.M., Lindor, N., Goode, E.L., Fridley, B.L., Jenkins, R., Le Marchand, L., Casey, G., Haile, R., Hopper, J., et al. 2010. Confirmation of linkage to and localization of familial colon cancer risk haplotype on chromosome 9q22. *Cancer Res.* 70: 5409-5418.

CHROMOSOMAL LOCATION

Genetic locus: ZNF169 (human) mapping to 9q22.32.

PRODUCT

ZNF169 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZNF169 shRNA Plasmid (h): sc-92480-SH and ZNF169 shRNA (h) Lentiviral Particles: sc-92480-V as alternate gene silencing products.

For independent verification of ZNF169 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-92480A, sc-92480B and sc-92480C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ZNF169 siRNA (h) is recommended for the inhibition of ZNF169 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZNF169 gene expression knockdown using RT-PCR Primer: ZNF169 (h)-PR: sc-92480-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.