



# CBE1 siRNA (h): sc-92698

## BACKGROUND

CBE1 (ciliated bronchial epithelium 1), also known as C9orf24 (chromosome 9 open reading frame 24) or testis development protein NYD-SP22, is a 262 amino acid protein that exists as three alternatively spliced isoforms. CBE1 is expressed in ciliated cells and is found at high levels in adult testis. Suggested to play a role in ciliogenesis and spermatogenesis, CBE1 is encoded by a gene that maps to human chromosome 9p13.3. Chromosome 9 consists of about 145 million bases, represents 4% of the human genome and encodes nearly 900 genes. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, and Familial dysautonomia, are both associated with chromosome 9. Notably, chromosome 9 encompasses the largest interferon family gene cluster.

## REFERENCES

1. Yoshisue, H., et al. 2004. Characterization of ciliated bronchial epithelium 1, a ciliated cell-associated gene induced during mucociliary differentiation. *Am. J. Respir. Cell Mol. Biol.* 31: 491-500.
2. Burmeister, T., et al. 2007. Atypical BCR-ABL mRNA transcripts in adult acute lymphoblastic leukemia. *Haematologica* 92: 1699-1702.
3. Cottin, V., et al. 2007. Pulmonary vascular manifestations of hereditary hemorrhagic telangiectasia (rendu-osler disease). *Respiration* 74: 361-378.
4. Zeitz, M.J., et al. 2009. Organization of the amplified type I interferon gene cluster and associated chromosome regions in the interphase nucleus of human osteosarcoma cells. *Chromosome Res.* 17: 305-319.
5. Haitchi, H.M., et al. 2009. Chronological expression of Ciliated Bronchial Epithelium 1 during pulmonary development. *Eur. Respir. J.* 33: 1095-1104.
6. Stassano, P., et al. 2010. Cardiac metastasis from malignant pancreatic somatostatinoma. *Surgery* 147: 172-174.

## CHROMOSOMAL LOCATION

Genetic locus: C9orf24 (human) mapping to 9p13.3.

## PRODUCT

CBE1 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see CBE1 shRNA Plasmid (h): sc-92698-SH and CBE1 shRNA (h) Lentiviral Particles: sc-92698-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

CBE1 siRNA (h) is recommended for the inhibition of CBE1 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor CBE1 gene expression knockdown using RT-PCR Primer: CBE1 (h)-PR: sc-92698-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.