



SURF-2 siRNA (h): sc-92780

BACKGROUND

SURF-2 (Surfeit locus protein 2) is a 256 amino acid protein that belongs to the SURF2 family and interacts with β -1,4-Gal-T3, uPAR and WDR20. The gene that encodes SURF-2 is located in the Surfeit gene cluster, which is a group of very tightly linked genes that do not share sequence similarity. The SURF-2 gene maps to human chromosome 9q34.2 and shares a bidirectional promoter with SURF1, which is located on the opposite strand. The intergenic region between the SURF-1 and SURF-2 genes is expected to have bidirectional promoter activity, as is found in mouse. Although this region lacks a TATA box, it is GC-rich. Housing over 900 genes, human chromosome 9 comprises nearly 4% of the human genome. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, and familial dysautonomia, are both associated with chromosome 9.

REFERENCES

1. Online Mendelian Inheritance in Man, OMIM™. 1989. Johns Hopkins University, Baltimore, MD. MIM Number: 185630. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
2. Yon, J., Jones, T., Garson, K., Sheer, D. and Fried, M. 1993. The organization and conservation of the human Surfeit gene cluster and its localization telomeric to the c-Abl and can proto-oncogenes at chromosome band 9q34.1. *Hum. Mol. Genet.* 2: 237-240.
3. Lennard, A., Gaston, K. and Fried, M. 1994. The Surf-1 and Surf-2 genes and their essential bidirectional promoter elements are conserved between mouse and human. *DNA Cell Biol.* 13: 1117-1126.
4. Duhig, T., Ruhrberg, C., Mor, O. and Fried, M. 1998. The human Surfeit locus. *Genomics* 52: 72-78.
5. Cottin, V., Dupuis-Girod, S., Lesca, G. and Cordier, J.F. 2007. Pulmonary vascular manifestations of hereditary hemorrhagic telangiectasia (Rendu-Osler disease). *Respiration* 74: 361-378.
6. Zeitz, M.J., Marella, N.V., Malyavantham, K.S., Goetze, S., Bode, J., Raska, I. and Berezney, R. 2009. Organization of the amplified type I interferon gene cluster and associated chromosome regions in the interphase nucleus of human osteosarcoma cells. *Chromosome Res.* 17: 305-319.
7. Gold-von Simson, G., Goldberg, J.D., Rolnitzky, L.M., Mull, J., Leyne, M., Voustianouk, A., Slangenaupt, S.A. and Axelrod, F.B. 2009. Kinetin in familial dysautonomia carriers: implications for a new therapeutic strategy targeting mRNA splicing. *Pediatr. Res.* 65: 341-346.
8. Axelrod, F.B., Hilz, M.J., Berlin, D., Yau, P.L., Javier, D., Sweat, V., Bruehl, H. and Convit, A. 2010. Neuroimaging supports central pathology in familial dysautonomia. *J. Neurol.* 257: 198-206.

CHROMOSOMAL LOCATION

Genetic locus: SURF2 (human) mapping to 9q34.2.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

SURF-2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see SURF-2 shRNA Plasmid (h): sc-92780-SH and SURF-2 shRNA (h) Lentiviral Particles: sc-92780-V as alternate gene silencing products.

For independent verification of SURF-2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-92780A, sc-92780B and sc-92780C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

SURF-2 siRNA (h) is recommended for the inhibition of SURF-2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor SURF-2 gene expression knockdown using RT-PCR Primer: SURF-2 (h)-PR: sc-92780-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.