# Metallothionein 1G siRNA (h): sc-93237



The Power to Question

#### **BACKGROUND**

Metallothionein (MT) is a sulfhydryl- and cysteine-rich protein found in microorganisms, plants and all invertebrate and vertebrate animals. Metallothioneins are a group of ubiquitous low-molecular-weight proteins that have functional roles in cell growth, repair and differentiation. They are implicated primarily in metal ion detoxification, in that they are essential for the protection of cells against the toxicity of cadmium, mercury and copper. Metallothionein, as an acute phase or stress-response protein and free radical scavenger, is related to inflammation and cellular protection from reactive forms of oxygen, ionizing radiation, pharmacological agents and mutagens. Metallothioneins are known to be broadly expressed in heart, liver, kidney, breast and testis tissue.

# **REFERENCES**

- loachim, E.E., et al. 2000. Immunohistochemical localization of Metallothionein in endometrial lesions. J. Pathol. 191: 269-273.
- Liu, J., et al. 2000. Metallothionein-I/II null mice are more sensitive than wild-type mice to the hepatotoxic and nephrotoxic effects of chronic oral or injected inorganic arsenicals. Toxicol. Sci. 55: 460-467.
- Cai, L., et al. 2000. Induction of Metallothionein synthesis with preservation of testicular function in rats following long term renal transplantation. Urol. Res. 28: 97-103.
- 4. Florianczyk, B., et al. 2000. Metallothionein levels in cell fractions from breast cancer tissues. Acta Oncol. 39: 141-143.
- 5. Theocharis, S.E., et al. 2000. Liver Metallothionein expression in thioacetamide-intoxicated rats. Pathol. Res. Pract. 196: 313-319.
- Kang, Y.J., et al. 2000. Metallothionein inhibits myocardial apoptosis in copper-deficient mice: role of atrial natriuretic peptide. Lab. Invest. 80: 745-757.
- Syring, R.A., et al. 2000. Cloning and sequencing of cDNAs encoding for a novel copper-specific Metallothionein and two cadmium-inducible Metallothioneins from the blue crab *Callinectes sapidus*. Comp. Biochem. Physiol. C Toxicol. Pharmacol. 125: 325-332.
- 8. Jayasurya, A., et al. 2000. Infiltrating lymphocytes in undifferentiated nasopharyngeal cancer lack Metallothionein expression. Cancer Lett. 155: 99-104.

# CHROMOSOMAL LOCATION

Genetic locus: MT1G (human) mapping to 16q13.

## **PRODUCT**

Metallothionein 1G siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu M$  solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Metallothionein 1G shRNA Plasmid (h): sc-93237-SH and Metallothionein 1G shRNA (h) Lentiviral Particles: sc-93237-V as alternate gene silencing products.

#### STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNAse-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNAse-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## **APPLICATIONS**

Metallothionein 1G siRNA (h) is recommended for the inhibition of Metallothionein 1G expression in human cells.

#### **SUPPORT REAGENTS**

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 µM in 66 µl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## **RESEARCH USE**

For research use only, not for use in diagnostic procedures.

## **PROTOCOLS**

See our web site at www.scbt.com for detailed protocols and support products.

**Santa Cruz Biotechnology, Inc.** 1.800.457.3801 831.457.3800 fax 831.457.3801 **Europe** +00800 4573 8000 49 6221 4503 0 **www.scbt.com**