



GPR97 siRNA (h): sc-93484

BACKGROUND

G protein-coupled receptors (GPRs), also known as seven transmembrane receptors, heptahelical receptors or 7TM receptors, comprise a superfamily of proteins that play a role in many different stimulus-response pathways. G protein-coupled receptors translate extracellular signals into intracellular signals (G protein activation) and they respond to a variety of signaling molecules, such as hormones and neurotransmitters. GPR97 (G protein-coupled receptor 97), also known as PB99 or PGR26, is a 549 amino acid multi-pass membrane protein that contains one GPS domain and belongs to the G protein-coupled receptor 2 family. GPR97 functions as an orphan receptor that is thought to play a role in signaling events throughout the cell.

REFERENCES

1. Larhammar, D., Blomqvist, A.G. and Wahlestedt, C. 1993. The receptor revolution—multiplicity of G protein-coupled receptors. *Drug Des. Discov.* 9: 179-188.
2. Ji, T.H., Grossmann, M. and Ji, I. 1998. G protein-coupled receptors. I. Diversity of receptor-ligand interactions. *J. Biol. Chem.* 273: 17299-17302.
3. Schöneberg, T., Schultz, G. and Gudermann, T. 1999. Structural basis of G protein-coupled receptor function. *Mol. Cell. Endocrinol.* 151: 181-193.
4. Fredriksson, R., Lagerström, M.C., Höglund, P.J. and Schiöth, H.B. 2002. Novel human G protein-coupled receptors with long N-terminals containing GPS domains and Ser/Thr-rich regions. *FEBS Lett.* 531: 407-414.
5. Vanti, W.B., Nguyen, T., Cheng, R., Lynch, K.R., George, S.R. and O'Dowd, B.F. 2003. Novel human G protein-coupled receptors. *Biochem. Biophys. Res. Commun.* 305: 67-71.
6. Vassilatis, D.K., Hohmann, J.G., Zeng, H., Li, F., Ranchalis, J.E., Mortrud, M.T., Brown, A., Rodriguez, S.S., Weller, J.R., Wright, A.C., Bergmann, J.E. and Gaitanaris, G.A. 2003. The G protein-coupled receptor repertoires of human and mouse. *Proc. Natl. Acad. Sci. USA* 100: 4903-4908.
7. Bjarnadóttir, T.K., Fredriksson, R., Höglund, P.J., Gloriam, D.E., Lagerström, M.C. and Schiöth, H.B. 2004. The human and mouse repertoire of the adhesion family of G protein-coupled receptors. *Genomics* 84: 23-33.
8. Bjarnadóttir, T.K., Geirardsdóttir, K., Ingemansson, M., Mirza, M.A., Fredriksson, R. and Schiöth, H.B. 2007. Identification of novel splice variants of Adhesion G protein-coupled receptors. *Gene* 387: 38-48.
9. Nordström, K.J., Lagerström, M.C., Waller, L.M., Fredriksson, R. and Schiöth, H.B. 2009. The Secretin GPCRs descended from the family of Adhesion GPCRs. *Mol. Biol. Evol.* 26: 71-84.

CHROMOSOMAL LOCATION

Genetic locus: ADGRG3 (human) mapping to 16q21.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

GPR97 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GPR97 shRNA Plasmid (h): sc-93484-SH and GPR97 shRNA (h) Lentiviral Particles: sc-93484-V as alternate gene silencing products.

For independent verification of GPR97 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-93484A, sc-93484B and sc-93484C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GPR97 siRNA (h) is recommended for the inhibition of GPR97 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GPR97 gene expression knockdown using RT-PCR Primer: GPR97 (h)-PR: sc-93484-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.