

β3Gn-T9 siRNA (h): sc-93513

BACKGROUND

A family of human β 1,3-galactosyltransferases (β3Gn-Ts) consists of nine members (β3Gn-T1, -T2, -T3, -T4, -T5, -T6, -T7, -T8 and -T9). β3Gn-T1 catalyzes the formation of type 1 oligosaccharides. β3Gn-T2 converts lacto-N-triose II into lacto-N-tetraose and lacto-N-neotetraose and can form a heterodimer with β3Gn-T8, which, as a complex, exhibits higher enzymatic activity. Unlike the ubiquitously expressed β3Gn-T2, β3Gn-T3 is specifically expressed in colon, jejunum, stomach, esophagus, placenta and trachea, and β3Gn-T4 is mainly expressed in brain. β3Gn-T5 is essential for the biosynthesis of Lewis antigens and may play a role in gastric cancer as a result of its participation in chronic *H. pylori* infection. β3Gn-T6 may be a useful marker for distinguishing between benign adenomas and premalignant lesions. β3Gn-T7 acts as an anti-migration factor for a lung cancer cell line. β3Gn-T9 is a single-pass type II membrane protein localized to the Golgi apparatus membrane.

REFERENCES

- Shiraishi, N., Natsume, A., Togayachi, A., Endo, T., Akashima, T., Yamada, Y., Imai, N., Nakagawa, S., Koizumi, S., Sekine, S., Narimatsu, H. and Sasaki, K. 2001. Identification and characterization of three novel β 1,3-N-acetylglucosaminyltransferases structurally related to the β 1,3-galactosyltransferase family. *J. Biol. Chem.* 276: 3498-3507.
- Seko, A. and Yamashita, K. 2004. β 1,3-N-acetylglucosaminyltransferase-7 (β3Gn-T7) acts efficiently on keratan sulfate-related glycans. *FEBS Lett.* 556: 216-220.
- Iwai, T., Kudo, T., Kawamoto, R., Kubota, T., Togayachi, A., Hiruma, T., Okada, T., Kawamoto, T., Morozumi, K. and Narimatsu, H. 2005. Core 3 synthase is down-regulated in colon carcinoma and profoundly suppresses the metastatic potential of carcinoma cells. *Proc. Natl. Acad. Sci. USA* 102: 4572-4577.
- Deo, V.K. and Park, E.Y. 2006. Multiple co-transfection and co-expression of human β 1,3-N-acetylglucosaminyltransferase with human calreticulin chaperone cDNA in a single step in insect cells. *Biotechnol. Appl. Biochem.* 43: 129-135.
- Seko, A. and Yamashita, K. 2008. Activation of β 1,3-N-acetylglucosaminyltransferase-2 (β3Gn-T2) by β3Gn-T8: possible involvement of β3Gn-T8 in increasing poly-N-acetyllactosamine chains in differentiated HL-60 cells. *J. Biol. Chem.* 283: 33094-33100.
- Marcos, N.T., Magalhães, A., Ferreira, B., Oliveira, M.J., Carvalho, A.S., Mendes, N., Gilmartin, T., Head, S.R., Figueiredo, C., David, L., Santos-Silva, F. and Reis, C.A. 2008. *Helicobacter pylori* induces β3Gn-T5 in human gastric cell lines, modulating expression of the SabA ligand sialyl-Lewis x. *J. Clin. Invest.* 118: 2325-2336.

CHROMOSOMAL LOCATION

Genetic locus: B3GNT9 (human) mapping to 16q22.1.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

β3Gn-T9 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μM solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see β3Gn-T9 shRNA Plasmid (h): sc-93513-SH and β3Gn-T9 shRNA (h) Lentiviral Particles: sc-93513-V as alternate gene silencing products.

For independent verification of β3Gn-T9 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-93513A, sc-93513B and sc-93513C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μl of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μl of RNase-free water makes a 10 μM solution in a 10 μM Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

β3Gn-T9 siRNA (h) is recommended for the inhibition of β3Gn-T9 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μM in 66 μl. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor β3Gn-T9 gene expression knockdown using RT-PCR Primer: β3Gn-T9 (h)-PR: sc-93513-PR (20 μl). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.