

# ZNF232 siRNA (h): sc-93736

## BACKGROUND

Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Krüppel-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins. ZNF232, also known as Zinc finger and SCAN domain-containing protein 11, is a 417 amino acid protein belonging to the Krüppel C<sub>2</sub>H<sub>2</sub>-type zinc-finger protein family. Localized to the nucleus, ZNF232 contains one SCAN box domain and five C<sub>2</sub>H<sub>2</sub>-type zinc fingers. Due to the presence of these domains, ZNF232 may be involved in transcriptional regulation. Ubiquitously expressed, ZNF232 is present at high levels in testis, liver and ovary. ZNF232 exists as two isoforms produced by alternative splicing.

## REFERENCES

1. Bellefroid, E.J., Poncelet, D.A., Lecocq, P.J., Revelant, O. and Martial, J.A. 1991. The evolutionarily conserved Krüppel-associated box domain defines a subfamily of eukaryotic multifingered proteins. *Proc. Natl. Acad. Sci. USA* 88: 3608-3612.
2. Pengue, G., Calabrò, V., Bartoli, P.C., Pagliuca, A. and Lania, L. 1994. Repression of transcriptional activity at a distance by the evolutionarily conserved KRAB domain present in a subfamily of zinc finger proteins. *Nucleic Acids Res.* 22: 2908-2914.
3. Margolin, J.F., Friedman, J.R., Meyer, W.K., Vissing, H., Thiesen, H.J. and Rauscher, F.J. 1994. Krüppel-associated boxes are potent transcriptional repression domains. *Proc. Natl. Acad. Sci. USA* 91: 4509-4513.
4. Williams, A.J., Blacklow, S.C. and Collins, T. 1999. The zinc finger-associated SCAN box is a conserved oligomerization domain. *Mol. Cell. Biol.* 19: 8526-8535.
5. Mavrogiannis, L.A., Argyrokastritis, A., Tzitzikas, N., Dermitzakis, E., Sarafidou, T., Patsalis, P.C. and Moschonas, N.K. 2001. ZNF232: structure and expression analysis of a novel human C<sub>2</sub>H<sub>2</sub> zinc finger gene, member of the SCAN/LeR domain subfamily. *Biochim. Biophys. Acta* 1518: 300-305.
6. Shannon, M., Hamilton, A.T., Gordon, L., Branscomb, E. and Stubbs, L. 2003. Differential expansion of zinc-finger transcription factor loci in homologous human and mouse gene clusters. *Genome Res.* 13: 1097-1110.

## CHROMOSOMAL LOCATION

Genetic locus: ZNF232 (human) mapping to 17p13.2.

## PRODUCT

ZNF232 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZNF232 shRNA Plasmid (h): sc-93736-SH and ZNF232 shRNA (h) Lentiviral Particles: sc-93736-V as alternate gene silencing products.

For independent verification of ZNF232 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-93736A, sc-93736B and sc-93736C.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

ZNF232 siRNA (h) is recommended for the inhibition of ZNF232 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZNF232 gene expression knockdown using RT-PCR Primer: ZNF232 (h)-PR: sc-93736-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.