

epsin 3 siRNA (h): sc-93806

BACKGROUND

Epsin 3 (EPN3), also known as EPS-15-interacting protein 3, is a 632 amino acid member of the epsin protein family. Epsin 3 localizes mainly to the perinuclear region of the nucleus and in clathrin-coated vesicles close to the cell periphery. Epsin 3 is expressed in migrating keratinocytes from wounded skin, but not in normal skin or differentiating keratinocytes. Epsin 3 contains one ENTH (epsin N-terminal homology) domain and two UIM (ubiquitin-interacting motif) repeats and is expressed as two isoforms produced by alternative splicing. The gene that encodes epsin 3 maps to human chromosome 17, which comprises over 2.5% of the human genome and encodes over 1,200 genes. Two key tumor suppressor genes are associated with chromosome 17, namely, p53 and BRCA1. Tumor suppressor p53 is necessary for maintenance of cellular genetic integrity by moderating cell fate through DNA repair versus cell death. Malfunction or loss of p53 expression is associated with malignant cell growth and Li-Fraumeni syndrome. Like p53, BRCA1 is directly involved in DNA repair, specifically it is recognized as a genetic determinant of early onset breast cancer and predisposition to cancers of the ovary, colon, prostate gland and fallopian tubes.

REFERENCES

1. Hall, J.M., Friedman, L., Guenther, C., Lee, M.K., Weber, J.L., Black, D.M. and King, M.C. 1992. Closing in on a breast cancer gene on chromosome 17q. *Am. J. Hum. Genet.* 50: 1235-1242.
2. Soussi, T., Dehouche, K. and Beroud, C. 2000. p53 website and analysis of p53 gene mutations in human cancer: forging a link between epidemiology and carcinogenesis. *Hum. Mutat.* 15: 105-113.
3. Spradling, K.D., Burke, T.J., Lohi, J. and Pilcher, B.K. 2000. Cloning and initial characterization of human epsin 3, a novel matrix-induced keratinocyte specific transcript. *J. Invest. Dermatol.* 115: 332.
4. Piura, B., Rabinovich, A. and Yanai-Inbar, I. 2001. Three primary malignancies related to BRCA mutation successively occurring in a BRCA1 185delAG mutation carrier. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 97: 241-244.
5. Spradling, K.D., McDaniel, A.E., Lohi, J. and Pilcher, B.K. 2001. Epsin 3 is a novel extracellular matrix-induced transcript specific to wounded epithelia. *J. Biol. Chem.* 276: 29257-29267.
6. Minamoto, T., Buschmann, T., Habelhah, H., Matusevich, E., Tahara, H., Boerresen-Dale, A.L., Harris, C., Sidransky, D. and Ronai, Z. 2001. Distinct pattern of p53 phosphorylation in human tumors. *Oncogene* 20: 3341-3347.
7. Colland, F., Jacq, X., Trouplin, V., Mougin, C., Groizeleau, C., Hamburger, A., Meil, A., Wojcik, J., Legrain, P. and Gauthier, J.M. 2004. Functional proteomics mapping of a human signaling pathway. *Genome Res.* 14: 1324-1332.

CHROMOSOMAL LOCATION

Genetic locus: EPN3 (human) mapping to 17q21.33.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

epsin 3 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see epsin 3 shRNA Plasmid (h): sc-93806-SH and epsin 3 shRNA (h) Lentiviral Particles: sc-93806-V as alternate gene silencing products.

For independent verification of epsin 3 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-93806A, sc-93806B and sc-93806C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

epsin 3 siRNA (h) is recommended for the inhibition of epsin 3 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor epsin 3 gene expression knockdown using RT-PCR Primer: epsin 3 (h)-PR: sc-93806-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.