

▶ TEX2 siRNA (h): sc-93815

BACKGROUND

TEX2 (testis-expressed sequence 2 protein) is a 1,127 amino acid multi-pass membrane protein that is specifically expressed in testis. There are two isoforms of TEX2 that are produced as a result of alternative splicing events. The gene encoding TEX2 maps to human chromosome 17q23.3, which comprises over 2.5% of the human genome and encodes over 1,200 genes. Two key tumor suppressor genes are associated with chromosome 17, namely, p53 and BRCA1. Tumor suppressor p53 is necessary for maintenance of cellular genetic integrity by moderating cell fate through DNA repair versus cell death. Malfunction or loss of p53 expression is associated with malignant cell growth and Li-Fraumeni syndrome. Chromosome 17 has also been implicated in the pathogenesis of frontotemporal dementia and parkinsonism.

REFERENCES

- James, C.D., et al. 1989. Mitotic recombination of chromosome 17 in astrocytomas. *Proc. Natl. Acad. Sci. USA* 86: 2858-2862.
- Lindblom, A., et al. 1993. Four separate regions on chromosome 17 show loss of heterozygosity in familial breast carcinomas. *Hum. Genet.* 91: 6-12.
- Negrini, M., et al. 1994. Tumor and growth suppression of breast cancer cells by chromosome 17-associated functions. *Cancer Res.* 54: 1818-1824.
- Clausen, O.P., et al. 1998. Association of p53 accumulation with TP53 mutations, loss of heterozygosity at 17p13, and DNA ploidy status in 273 colorectal carcinomas. *Diagn. Mol. Pathol.* 7: 215-223.
- Gilbert, F. 1998. Disease genes and chromosomes: disease maps of the human genome. *Chromosome 17. Genet. Test.* 2: 357-381.
- Risio, M., et al. 2003. Deletions of 17p are associated with transition from early to advanced colorectal cancer. *Cancer Genet. Cytogenet.* 147: 44-49.
- Wszolek, Z.K., et al. 2006. Frontotemporal dementia and parkinsonism linked to chromosome 17 (FTDP-17). *Orphanet J. Rare Dis.* 1: 30.
- Mayya, V., et al. 2009. Quantitative phosphoproteomic analysis of T cell receptor signaling reveals system-wide modulation of protein-protein interactions. *Sci. Signal.* 2: ra46.

CHROMOSOMAL LOCATION

Genetic locus: TEX2 (human) mapping to 17q23.3.

PRODUCT

TEX2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see TEX2 shRNA Plasmid (h): sc-93815-SH and TEX2 shRNA (h) Lentiviral Particles: sc-93815-V as alternate gene silencing products.

For independent verification of TEX2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-93815A, sc-93815B and sc-93815C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

TEX2 siRNA (h) is recommended for the inhibition of TEX2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor TEX2 gene expression knockdown using RT-PCR Primer: TEX2 (h)-PR: sc-93815-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.