

ZBP2 siRNA (h): sc-93858

BACKGROUND

Mammalian gametes recognize each other when the plasma membrane of the sperm head binds to the zona pellucida (ZP), an extracellular coat surrounding eggs. The zona pellucida is composed of four major glycoproteins expressed in oocytes, ZP1, ZP2, ZP3 and ZP4. Important receptors of zona pellucida on sperm are the zona pellucida binding proteins (ZPBPs). ZPBPs are suggested to play an essential role in sperm-egg interaction. ZBP2 (zona pellucida binding protein 2), also known as ZPBPL, is a 338 amino acid secreted protein that belongs to the zona pellucida-binding protein Sp38 family. Mutations in the gene encoding ZBP2 may cause subfertility. ZBP2 exists as two alternatively spliced isoforms and is encoded by a gene located on human chromosome 17q12.

REFERENCES

1. Liang, L.F., Chamow, S.M. and Dean, J. 1990. Oocyte-specific expression of mouse Zp-2: developmental regulation of the zona pellucida genes. *Mol. Cell. Biol.* 10: 1507-1515.
2. Dean, J. 1992. Biology of mammalian fertilization: role of the zona pellucida. *J. Clin. Invest.* 89: 1055-1059.
3. Kipersztok, S., Osawa, G.A., Liang, L.F., Modi, W.S. and Dean, J. 1995. POM-ZP3, a bipartite transcript derived from human ZP3 and a POM121 homologue. *Genomics* 25: 354-359.
4. Katoh, M. and Katoh, M. 2003. Identification and characterization of human ZBP2-like gene in silico. *Int. J. Mol. Med.* 12: 399-404.
5. Jazwinska, A. and Affolter, M. 2004. A family of genes encoding zona pellucida (ZP) domain proteins is expressed in various epithelial tissues during *Drosophila* embryogenesis. *Gene Expr. Patterns* 4: 413-421.
6. Wassarman, P.M., Jovine, L., Litscher, E.S., Qi, H. and Williams, Z. 2004. Egg-sperm interactions at fertilization in mammals. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 115: S57-S60.
7. Lin, Y.N., Roy, A., Yan, W., Burns, K.H. and Matzuk, M.M. 2007. Loss of zona pellucida binding proteins in the acrosomal matrix disrupts acrosome biogenesis and sperm morphogenesis. *Mol. Cell. Biol.* 27: 6794-6805.
8. Verlaan, D.J., Berlivet, S., Hunninghake, G.M., Madore, A.M., Larivière, M., Moussette, S., Grundberg, E., Kwan, T., Ouimet, M., Ge, B., Hoberman, R., Swiatek, M., Dias, J., Lam, K.C., Koka, V., Harmsen, E., et al. 2009. Allele-specific chromatin remodeling in the ZBP2/GSDMB/ORMDL3 locus associated with the risk of asthma and autoimmune disease. *Am. J. Hum. Genet.* 85: 377-393.
9. Song, C., Zhou, H., Gao, B., Sun, L., Wu, H., Wang, X., Chen, G. and Mao, J. 2010. Molecular cloning of pig ZBP2 and mRNA expression of ZBP1 and ZBP2 in reproductive tracts of boars. *Anim. Reprod. Sci.* 122: 229-235.

CHROMOSOMAL LOCATION

Genetic locus: ZBP2 (human) mapping to 17q12.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

ZBP2 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see ZBP2 shRNA Plasmid (h): sc-93858-SH and ZBP2 shRNA (h) Lentiviral Particles: sc-93858-V as alternate gene silencing products.

For independent verification of ZBP2 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-93858A, sc-93858B and sc-93858C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

ZBP2 siRNA (h) is recommended for the inhibition of ZBP2 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor ZBP2 gene expression knockdown using RT-PCR Primer: ZBP2 (h)-PR: sc-93858-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.