

# Sidekick-2 siRNA (h): sc-94033

## BACKGROUND

Sidekick-2 (SDK2), also known as KIAA1514, is a 2,172 amino acid single-pass type I membrane protein containing thirteen fibronectin type-III domains and six Ig-like C2-type (immunoglobulin-like) domains. Acting as a cell adhesion protein, Sidekick-2 guides axonal terminals to synapses in developing neurons. Sidekick-2 may require the interaction with MAGI scaffolding proteins for proper localization and function. Existing as four alternatively spliced isoforms, the gene encoding Sidekick-2 maps to human chromosome 17q25.1. Chromosome 17 makes up over 2.5% of the human genome with about 81 million bases encoding over 1,200 genes. Neurofibromatosis, a condition characterized by neural and epidermal lesions, and dysregulated Schwann cell growth have been linked to chromosome 17. Alexander disease, Birt-Hogg-Dube syndrome and Canavan disease are also associated with chromosome 17.

## REFERENCES

1. Yamagata, M., Weiner, J.A. and Sanes, J.R. 2002. Sidekicks: synaptic adhesion molecules that promote lamina-specific connectivity in the retina. *Cell* 110: 649-660.
2. Kaufman, L., Hayashi, K., Ross, M.J., Ross, M.D. and Klotman, P.E. 2004. Sidekick-1 is upregulated in glomeruli in HIV-associated nephropathy. *J. Am. Soc. Nephrol.* 15: 1721-1730.
3. Hayashi, K., Kaufman, L., Ross, M.D. and Klotman, P.E. 2005. Definition of the critical domains required for homophilic targeting of mouse sidekick molecules. *FASEB J.* 19: 614-616.
4. Zody, M.C., Garber, M., Adams, D.J., Sharpe, T., Harrow, J., Lupski, J.R., Nicholson, C., Searle, S.M., Wilming, L., Young, S.K., Abouelleil, A., Allen, N.R., Bi, W., Bloom, T., Borowsky, M.L., Bugalter, B.E., Butler, J., et al. 2006. DNA sequence of human chromosome 17 and analysis of rearrangement in the human lineage. *Nature* 440: 1045-1049.
5. Yamagata, M. and Sanes, J.R. 2008. Dscam and Sidekick proteins direct lamina-specific synaptic connections in vertebrate retina. *Nature* 451: 465-469.
6. Gauci, S., Helbig, A.O., Slijper, M., Krijgsveld, J., Heck, A.J. and Mohammed, S. 2009. Lys-N and trypsin cover complementary parts of the phosphoproteome in a refined SCX-based approach. *Anal. Chem.* 81: 4493-4501.
7. Yamagata, M. and Sanes, J.R. 2010. Synaptic localization and function of Sidekick recognition molecules require MAGI scaffolding proteins. *J. Neurosci.* 30: 3579-3588.

## CHROMOSOMAL LOCATION

Genetic locus: SDK2 (human) mapping to 17q25.1.

## PROTOCOLS

See our web site at [www.scbt.com](http://www.scbt.com) for detailed protocols and support products.

## PRODUCT

Sidekick-2 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10  $\mu$ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Sidekick-2 shRNA Plasmid (h): sc-94033-SH and Sidekick-2 shRNA (h) Lentiviral Particles: sc-94033-V as alternate gene silencing products.

## STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330  $\mu$ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330  $\mu$ l of RNase-free water makes a 10  $\mu$ M solution in a 10  $\mu$ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

## APPLICATIONS

Sidekick-2 siRNA (h) is recommended for the inhibition of Sidekick-2 expression in human cells.

## SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10  $\mu$ M in 66  $\mu$ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

## RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Sidekick-2 gene expression knockdown using RT-PCR Primer: Sidekick-2 (h)-PR: sc-94033-PR (20  $\mu$ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

## RESEARCH USE

For research use only, not for use in diagnostic procedures.