



Ribosomal Protein L27 siRNA (h): sc-94077

BACKGROUND

Ribosomes, the organelles that catalyze protein synthesis, are composed of a small subunit (40S) and a large subunit (60S) that consist of over 80 distinct ribosomal proteins. Mammalian ribosomal proteins are encoded by multigene families that contain processed pseudogenes and one functional intron-containing gene within their coding regions. Ribosomal Protein L27, also known as RPL27, is a 136 amino acid protein belonging to the Ribosomal Protein L27e family exists as a component of the 60S subunit, possibly playing a role in protein translation. Like most ribosomal proteins, Ribosomal Protein L27 exists as multiple processed pseudogenes that are scattered throughout the genome. Considered a novel candidate housekeeping gene, the gene encoding Ribosomal Protein L27 maps to human chromosome 17, which comprises over 2.5% of the human genome and encodes over 1,200 genes.

REFERENCES

1. Gallagher, R.A., et al. 1994. Cloning and nucleotide sequence of a full length cDNA encoding Ribosomal Protein L27 from human fetal kidney. *Biochim. Biophys. Acta* 1217: 329-332.
2. Fujiwara, S., et al. 1994. The gene for Ribosomal Protein L27 is located on the plastid rather than the nuclear genome of the chlorophyll c-containing alga *Pleurochrysis carterae*. *Plant Mol. Biol.* 24: 253-257.
3. Wool, I.G., et al. 1995. Structure and evolution of mammalian ribosomal proteins. *Biochem. Cell Biol.* 73: 933-947.
4. Varis, A., et al. 2002. Targets of gene amplification and overexpression at 17q in gastric cancer. *Cancer Res.* 62: 2625-2629.
5. Online Mendelian Inheritance in Man, OMIM™. 2003. Johns Hopkins University, Baltimore, MD. MIM Number: 607526. World Wide Web URL: <http://www.ncbi.nlm.nih.gov/omim/>
6. Ueda, M., et al. 2006. Promoter shuffling at a nuclear gene for mitochondrial RPL27. Involvement of interchromosome and subsequent intrachromosome recombinations. *Plant Physiol.* 141: 702-710.
7. Silva, E., et al. 2009. Prostaglandin synthesis genes are differentially transcribed in normal and pyometra endometria of bitches. *Reprod. Domest. Anim.* 44: 200-203.

CHROMOSOMAL LOCATION

Genetic locus: RPL27 (human) mapping to 17q21.31.

PRODUCT

Ribosomal Protein L27 siRNA (h) is a target-specific 19-25 nt siRNA designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see Ribosomal Protein L27 shRNA Plasmid (h): sc-94077-SH and Ribosomal Protein L27 shRNA (h) Lentiviral Particles: sc-94077-V as alternate gene silencing products.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

Ribosomal Protein L27 siRNA (h) is recommended for the inhibition of Ribosomal Protein L27 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor Ribosomal Protein L27 gene expression knockdown using RT-PCR Primer: Ribosomal Protein L27 (h)-PR: sc-94077-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.