

KIF19 siRNA (h): sc-94096

BACKGROUND

The kinesins constitute a large family of microtubule-dependent motor proteins, which are responsible for the distribution of numerous organelles, vesicles and macromolecular complexes throughout the cell. Kinesins also play crucial roles in cell division, intracellular transport and membrane trafficking events including endocytosis and transcytosis. Belonging to the Kinesin-like protein family, KIF19 is a 998 amino acid protein that contains one kinesin-motor domain, a domain that creates force in order to bind and move on microtubules. The gene encoding KIF19 maps to human chromosome 17, which comprises over 2.5% of the human genome and encodes over 1,200 genes. There are three isoforms of KIF19 that are produced as a result of alternative splicing events.

REFERENCES

1. Vallee, R.B. and Shpetner, H.S. 1990. Motor proteins of cytoplasmic microtubules. *Annu. Rev. Biochem.* 59: 909-932.
2. Endow, S.A. 1991. The emerging kinesin family of microtubule motor proteins. *Trends Biochem. Sci.* 16: 221-225.
3. Bloom, G.S. and Endow, S.A. 1995. Motor proteins 1: kinesins. *Protein Profile* 2: 1105-1171.
4. Brady, S.T. 1995. A kinesin medley: biochemical and functional heterogeneity. *Trends Cell Biol.* 5: 159-164.
5. Sablin, E.P., Kull, F.J., Cooke, R., Vale, R.D. and Fletterick, R.J. 1996. Crystal structure of the motor domain of the kinesin-related motor ncd. *Nature* 380: 555-559.
6. Kozielski, F., Sack, S., Marx, A., Thormählen, M., Schönbrunn, E., Biou, V., Thompson, A., Mandelkow, E.M. and Mandelkow, E. 1997. The crystal structure of dimeric kinesin and implications for microtubule-dependent motility. *Cell* 91: 985-994.
7. Vinogradova, M.V., Malanina, G.G., Reddy, V.S., Reddy, A.S. and Fletterick, R.J. 2008. Structural dynamics of the microtubule binding and regulatory elements in the kinesin-like calmodulin binding protein. *J. Struct. Biol.* 163: 76-83.
8. Wang, X. and Schwarz, T.L. 2009. The mechanism of Ca²⁺-dependent regulation of kinesin-mediated mitochondrial motility. *Cell* 136: 163-174.

CHROMOSOMAL LOCATION

Genetic locus: KIF19 (human) mapping to 17q25.1.

PRODUCT

KIF19 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see KIF19 shRNA Plasmid (h): sc-94096-SH and KIF19 shRNA (h) Lentiviral Particles: sc-94096-V as alternate gene silencing products.

For independent verification of KIF19 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-94096A, sc-94096B and sc-94096C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNases and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

KIF19 siRNA (h) is recommended for the inhibition of KIF19 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor KIF19 gene expression knockdown using RT-PCR Primer: KIF19 (h)-PR: sc-94096-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.