

GPR179 siRNA (h): sc-94166

BACKGROUND

G protein-coupled receptors (GPRs), also known as seven transmembrane receptors, heptahelical receptors or 7TM receptors, comprise a superfamily of proteins that play a role in many different stimulus-response pathways. G protein-coupled receptors translate extracellular signals into intracellular signals (G protein activation) and they respond to a variety of signaling molecules, such as hormones and neurotransmitters. GPR179, also known as GPR158L1 (G protein-coupled receptor 158-like 1), is a 2,367 amino acid multi-pass membrane protein that functions as an orphan receptor and belongs to the GPR3 family. The gene encoding GPR179 maps to human chromosome 17, which comprises over 2.5% of the human genome and encodes over 1,200 genes. Two key tumor suppressor genes are associated with chromosome 17, namely, p53 and BRCA1.

REFERENCES

1. Soussi, T., Dehouche, K. and Beroud, C. 2000. p53 website and analysis of p53 gene mutations in human cancer: forging a link between epidemiology and carcinogenesis. *Hum. Mutat.* 15: 105-113.
2. Piura, B., Rabinovich, A. and Yanai-Inbar, I. 2001. Three primary malignancies related to BRCA mutation successively occurring in a BRCA1 185delAG mutation carrier. *Eur. J. Obstet. Gynecol. Reprod. Biol.* 97: 241-244.
3. Minamoto, T., Buschmann, T., Habelhah, H., Matusевич, E., Tahara, H., Boerresen-Dale, A.L., Harris, C., Sidransky, D. and Ronai, Z. 2001. Distinct pattern of p53 phosphorylation in human tumors. *Oncogene* 20: 3341-3347.
4. Menzaghi, F., Behan, D.P. and Chalmers, D.T. 2002. Constitutively activated G protein-coupled receptors: a novel approach to CNS drug discovery. *Curr. Drug. Targets CNS Neurol. Disord.* 1: 105-121.
5. Szekeres, P.G. 2002. Functional assays for identifying ligands at orphan G protein-coupled receptors. *Recept. Channels* 8: 297-308.
6. Vassilatis, D.K., Hohmann, J.G., Zeng, H., Li, F., Ranchalis, J.E., Mortrud, M.T., Brown, A., Rodriguez, S.S., Weller, J.R., Wright, A.C., Bergmann, J.E. and Gaitanaris, G.A. 2003. The G protein-coupled receptor repertoires of human and mouse. *Proc. Natl. Acad. Sci. USA* 100: 4903-4908.
7. Bjarnadóttir, T.K., Fredriksson, R. and Schiöth, H.B. 2005. The gene repertoire and the common evolutionary history of glutamate, pheromone (V2R), taste(1) and other related G protein-coupled receptors. *Gene* 362: 70-84.
8. Davies, M.N., Gloriam, D.E., Secker, A., Freitas, A.A., Timmis, J. and Flower, D.R. 2011. Present perspectives on the automated classification of the G protein-coupled receptors (GPCRs) at the protein sequence level. *Curr. Top. Med. Chem.* 11: 1994-2009.

CHROMOSOMAL LOCATION

Genetic locus: GPR179 (human) mapping to 17q12.

PROTOCOLS

See our web site at www.scbt.com for detailed protocols and support products.

PRODUCT

GPR179 siRNA (h) is a pool of 3 target-specific 19-25 nt siRNAs designed to knock down gene expression. Each vial contains 3.3 nmol of lyophilized siRNA, sufficient for a 10 μ M solution once resuspended using protocol below. Suitable for 50-100 transfections. Also see GPR179 shRNA Plasmid (h): sc-94166-SH and GPR179 shRNA (h) Lentiviral Particles: sc-94166-V as alternate gene silencing products.

For independent verification of GPR179 (h) gene silencing results, we also provide the individual siRNA duplex components. Each is available as 3.3 nmol of lyophilized siRNA. These include: sc-94166A, sc-94166B and sc-94166C.

STORAGE AND RESUSPENSION

Store lyophilized siRNA duplex at -20° C with desiccant. Stable for at least one year from the date of shipment. Once resuspended, store at -20° C, avoid contact with RNAses and repeated freeze thaw cycles.

Resuspend lyophilized siRNA duplex in 330 μ l of the RNase-free water provided. Resuspension of the siRNA duplex in 330 μ l of RNase-free water makes a 10 μ M solution in a 10 μ M Tris-HCl, pH 8.0, 20 mM NaCl, 1 mM EDTA buffered solution.

APPLICATIONS

GPR179 siRNA (h) is recommended for the inhibition of GPR179 expression in human cells.

SUPPORT REAGENTS

For optimal siRNA transfection efficiency, Santa Cruz Biotechnology's siRNA Transfection Reagent: sc-29528 (0.3 ml), siRNA Transfection Medium: sc-36868 (20 ml) and siRNA Dilution Buffer: sc-29527 (1.5 ml) are recommended. Control siRNAs or Fluorescein Conjugated Control siRNAs are available as 10 μ M in 66 μ l. Each contain a scrambled sequence that will not lead to the specific degradation of any known cellular mRNA. Fluorescein Conjugated Control siRNAs include: sc-36869, sc-44239, sc-44240 and sc-44241. Control siRNAs include: sc-37007, sc-44230, sc-44231, sc-44232, sc-44233, sc-44234, sc-44235, sc-44236, sc-44237 and sc-44238.

RT-PCR REAGENTS

Semi-quantitative RT-PCR may be performed to monitor GPR179 gene expression knockdown using RT-PCR Primer: GPR179 (h)-PR: sc-94166-PR (20 μ l). Annealing temperature for the primers should be 55-60° C and the extension temperature should be 68-72° C.

RESEARCH USE

For research use only, not for use in diagnostic procedures.